



MMWRTM

Morbidity and Mortality Weekly Report

Weekly

November 22, 2002 / Vol. 51 / No. 46

Increases in Fluoroquinolone-Resistant *Neisseria gonorrhoeae* — Hawaii and California, 2001


Neisseria gonorrhoeae is a major cause of pelvic inflammatory disease, ectopic pregnancy, and infertility, and it can facilitate human immunodeficiency virus (HIV) transmission (1). Gonorrhea is the second most frequently reported communicable disease in the United States, with 361,705 reported cases in 2001 (2). During the 1980s, gonococcal resistance to penicillin and tetracycline became widespread; as a result, CDC recommended using cephalosporins as first-line treatment for gonorrhea. Since 1993, CDC also has recommended using fluoroquinolones (i.e., ciprofloxacin, ofloxacin, or levofloxacin) for gonorrhea treatment. Fluoroquinolone therapy is used widely because it is a relatively inexpensive, oral, and single-dose therapy. However, fluoroquinolone-resistant *N. gonorrhoeae* (QRNG)* is being identified more frequently (3). This report summarizes investigations of increases in QRNG in Hawaii and California in 2001 and provides data to support the recommendation that cephalosporins (i.e., ceftriaxone or cefixime) be used instead of fluoroquinolones as first-line treatment for gonorrhea acquired in these two states. The increases in QRNG highlight the importance of monitoring gonococcal resistance throughout the United States to guide local treatment decisions.

Hawaii

In 2001, the Hawaii State Laboratory performed gonorrhea culture and antimicrobial susceptibility tests on specimens from 265 (44%) of 605 reported gonorrhea cases. Patients seeking care at the public sexually transmitted disease (STD) clinic accounted for 44% (117 of 265) of these isolates. Overall, QRNG accounted for 20% (53 of 265) of

gonococcal isolates tested, compared with 11% in 2000 and 10% in 1999 (Figure 1). In 2001, 36% (19 of 53) of QRNG infections were among STD clinic patients.

FIGURE 1. Percentage of fluoroquinolone-resistant *Neisseria gonorrhoeae* (QRNG)* among tested gonococcal isolates and gonorrhea rate†, by year‡ — Hawaii, 1993–2001



* Defined as *N. gonorrhoeae* resistant to ciprofloxacin (minimal inhibitory concentration [MIC] ≥ 1.0 $\mu\text{g/mL}$ by agar dilution or disk diffusion zone size ≤ 27 mm) or ofloxacin (MIC ≥ 2.0 $\mu\text{g/mL}$ or disk diffusion zone size ≤ 24 mm) by the National Committee on Clinical Laboratory Standards.

† Per 100,000 population.

‡ Data for 1993–2001 include Gonococcal Isolate Surveillance Project (GISP) and non-GISP isolates.

INSIDE

- 1044 Multistate Outbreaks of *Salmonella* Serotype Poona Infections Associated with Eating Cantaloupe from Mexico — United States and Canada, 2000–2002
- 1047 Enterovirus Surveillance — United States, 2000–2001
- 1049 West Nile Virus Activity — United States, November 14–20, 2002, and Missouri, January 1–November 9, 2002
- 1051 Approval of a New Rapid Test for HIV Antibody
- 1052 Discontinuation of Cefixime Tablets — United States

* Defined as *N. gonorrhoeae* resistant to ciprofloxacin (minimal inhibitory concentration [MIC] ≥ 1.0 $\mu\text{g/mL}$ by agar dilution or disk diffusion zone size ≤ 27 mm) or ofloxacin (MIC ≥ 2.0 $\mu\text{g/mL}$ or disk diffusion zone size ≤ 24 mm) by the National Committee on Clinical Laboratory Standards.

CENTERS FOR DISEASE CONTROL AND PREVENTION

SAFER • HEALTHIER • PEOPLETM

The *MMWR* series of publications is published by the Epidemiology Program Office, Centers for Disease Control and Prevention (CDC), U.S. Department of Health and Human Services, Atlanta, GA 30333.

SUGGESTED CITATION

Centers for Disease Control and Prevention. [Article Title]. *MMWR* 2002;51:[inclusive page numbers].

Centers for Disease Control and Prevention

Julie L. Gerberding, M.D., M.P.H.

Director

David W. Fleming, M.D.

Deputy Director for Science and Public Health

Dixie E. Snider, Jr., M.D., M.P.H.

Associate Director for Science

Epidemiology Program Office

Stephen B. Thacker, M.D., M.Sc.

Director

Office of Scientific and Health Communications

John W. Ward, M.D.

Director

Editor, MMWR Series

David C. Johnson

Acting Managing Editor, MMWR (Weekly)

Jude C. Rutledge

Teresa F. Rutledge

Jeffrey D. Sokolow, M.A.

Writers/Editors, MMWR (Weekly)

Lynda G. Cupell

Malbea A. Heilman

Beverly J. Holland

Visual Information Specialists

Quang M. Doan

Erica R. Shaver

Information Technology Specialists

Division of Public Health Surveillance and Informatics

Notifiable Disease Morbidity and 122 Cities Mortality Data

Robert F. Fagan

Deborah A. Adams

Felicia J. Connor

Lateka Dammond

Patsy A. Hall

Pearl C. Sharp


Medical and interview records of the 117 STD clinic patients with positive gonococcal cultures diagnosed during January–December 2001 were reviewed to identify risk factors for QRNG; 19 (16%) had QRNG isolates. QRNG prevalence was higher for men who had sex exclusively with women than for men who had sex with men (MSM) (11 [20%] of 55 versus one [3%] of 29; $p=0.05$). Persons with a history of recent travel to Asia or a sex partner with such a history were not significantly more likely to have QRNG (four [36%] of 11) than persons without such a history (14 [14%] of 102; $p=0.07$). Unlike in Hawaii in 1999 (4), QRNG prevalence was not significantly higher among Asians/Pacific Islanders than among non-Asians/Pacific Islanders (10 [19%] of 54 versus nine [14%] of 63; $p=0.54$).

Since 2000, the Hawaii Department of Health (HDH) has recommended that clinicians avoid using fluoroquinolones to treat gonorrhea. Because of the 25% increase in reported gonorrhea morbidity (from 39.9 cases per 100,000 population in 2000 to 49.9 in 2001), adherence to this recommendation is particularly important. In February 2002, HDH informed all clinicians of the increases in gonorrhea and QRNG and organized STD training for an expanded network of clinicians and workers in community-based organizations. Preliminary analysis of gonococcal susceptibility results for 147 patients during January–June 2002 suggests that QRNG prevalence remains >14%.

California

San Francisco, Long Beach, Orange County, and San Diego are participants in the Gonococcal Isolate Surveillance Project (GISP), a CDC-sponsored sentinel surveillance system that monitors antimicrobial resistance in *N. gonorrhoeae* through antimicrobial susceptibility testing of male urethral gonococcal isolates obtained from patients at public STD clinics in 26 U.S. cities. During 1990–2000, <1% of isolates tested annually from each GISP site in California were QRNG, except for Orange County, where 5.6% (six of 107) of GISP isolates were QRNG in 2000. In 2001, susceptibility testing was expanded beyond the GISP sample to include all gonococcal isolates from Orange County and San Diego STD clinic patients, including those from women and nonurethral sites. Susceptibility testing also was performed on all gonococcal isolates obtained from patients at a large southern California health maintenance organization (HMO) during February–April 2001. In 2001, QRNG was identified in 2.5% (33 of 1,311) of patients with tested isolates (Figure 2). Among STD clinic patients with gonorrhea, 3.4% (10 of 297) in San Francisco, 3.0% (three of 99) in Long Beach, 3.3% (seven of 212) in Orange County, and 2.4% (eight of 330) in San Diego had QRNG. Among HMO patients with gonorrhea, 1.3% (five

FIGURE 2. Percentage of fluoroquinolone-resistant *Neisseria gonorrhoeae* (QRNG)* among tested gonococcal isolates and gonorrhea rate†, by year§ — California, 1993–2001



* Defined as *N. gonorrhoeae* resistant to ciprofloxacin (minimal inhibitory concentration [MIC] ≥ 1.0 $\mu\text{g/mL}$ by agar dilution or disk diffusion zone size ≤ 27 mm) or ofloxacin (MIC ≥ 2.0 $\mu\text{g/mL}$ or disk diffusion zone size ≤ 24 mm) by the National Committee on Clinical Laboratory Standards.

† Per 100,000 population.

§ Data for 1993–2000 are exclusively from the Gonococcal Isolate Surveillance Project (GISP), and data for 2001 include GISP and non-GISP isolates.

of 373) had QRNG. The 1,311 patients with tested isolates accounted for 5.6% of all reported gonorrhea cases in California in 2001. Among 29 men infected with QRNG in 2001 whose sexual orientation was known, 20 (69%) were MSM. Among MSM with QRNG, 19 had a median of three recent (within 2–6 months) sex partners (range: one–40); 10 heterosexual men and women with QRNG had a median of 1.5 recent sex partners (range: one–eight), indicating the potential for more rapid spread among MSM. Although 12 (43%) of 28 QRNG patients interviewed in 2001 reported recent travel to Asia, the Pacific Islands, or Hawaii by themselves or a sex partner, 57% denied such travel, suggesting endemic spread of QRNG within California.

Medical records were reviewed for all 469 gonorrhea patients whose isolates were tested for susceptibility and who were seen in San Francisco, Long Beach, Orange County, or San Diego STD clinics during July 1–December 31, 2001. QRNG was identified in 23 (4.9%) of the 469 patients tested. QRNG was more common among Asians/Pacific Islanders than among non-Asians/Pacific Islanders (four [16.7%] of 24 versus 19 [4.4%] of 427; $p=0.03$). QRNG prevalence was similar among MSM (5.2% [15 of 289]), heterosexual men (4.7% [seven of 149]), and women (3.6% [one of 28]). However, geographic variation was noted in California: in San Diego, QRNG was more common among MSM than among heterosexual men and women (6.6% [seven of 106] versus zero of 65; $p=0.03$), and in San Francisco, QRNG was more common among heterosexuals than among MSM (11.4% [five of 44] versus 1.1% [one of 93]; $p=0.01$).

In response to the increasing prevalence of QRNG, in May 2002 the California Department of Health Services advised clinicians to avoid using fluoroquinolones for treatment of gonorrhea. Preliminary data collected during January–June 2002 indicate that the prevalence of QRNG infection among STD clinic patients with tested gonococcal isolates in GISP sites in California has increased, exceeding 9% during this period.

Reported by: PM Whiticar, RG Ohye, MS, MV Lee, MS, Hawaii State Dept of Health. HM Bauer, MD, G Bolan, MD, STD Control Br, California Dept of Health Svcs. SA Wang, MD, RA Gunn, MD, HS Weinstock, MD, SM Berman, MD, Div of STD Prevention, National Center for HIV, STD, and TB Prevention; KE Mark, MD, LM Newman, MD, EIS officers, CDC.

Editorial Note: These data demonstrate that in 2001, QRNG prevalence increased in Hawaii and in California, where the epidemiology of QRNG varies within the state. In California, antimicrobial susceptibility data are available for a smaller proportion of reported gonorrhea cases than in Hawaii (6% versus 44%). Demographic data suggest that this low proportion might limit the generalizability of California's findings: patients with susceptibility-tested isolates in California were more likely to be male, older, and white, and to have their condition diagnosed in STD clinics than were other gonorrhea patients. However, the data from California indicate that QRNG has reached the continental United States, increasing the risk for its spread. Sporadic cases of QRNG have been identified in other states through GISP and non-GISP reporting, but no sustained increase in QRNG $>1\%$ has been identified in any other state (3). Increases in QRNG in California and Hawaii highlight the ongoing need for monitoring antimicrobial susceptibilities of gonococcal isolates throughout the United States.

CDC recommends that fluoroquinolones not be used to treat gonococcal infections acquired in Asia, where QRNG prevalence exceeds 40% (5); in the Pacific Islands, including Hawaii; in California; and in other areas with increased prevalence of fluoroquinolone resistance (6). The recommended treatment options for persons who might have acquired infection in those areas are cefixime (7), ceftriaxone, or spectinomycin. To select appropriate gonorrhea treatment in areas outside Hawaii and California, clinicians should ask suspected gonorrhea patients about their recent travel history and that of their sex partners (8).

Treatment of gonorrhea with fluoroquinolones can continue in areas where the prevalence of resistance is $<1\%$ (9). In areas where resistance is $\geq 1\%$, health departments making local treatment recommendations for gonorrhea also should consider other local factors such as the overall prevalence of

gonorrhea, the availability of antimicrobial susceptibility data, and the cost of various diagnostic and treatment options (10). Fluoroquinolones remain an important gonorrhea treatment option in the United States because they are inexpensive and easy to administer. In addition, their use might decrease use of cephalosporins and delay the development of cephalosporin resistance.

As part of effective gonorrhea control, state health departments should monitor local gonococcal antimicrobial susceptibility prevalence routinely to assist in developing local treatment recommendations. Symptomatic treatment failures are not a reliable indicator of emerging antimicrobial resistance because gonococcal infections, especially in women, are frequently asymptomatic. In 2001, a survey of STD project areas found that nonculture gonococcal tests were used widely and that approximately half of project areas had antimicrobial susceptibility data (3). Because nonculture tests cannot provide antimicrobial susceptibility results, local gonococcal culture capacity should be maintained. The antimicrobial susceptibility testing panel should, at a minimum, include a fluoroquinolone, cefixime, ceftriaxone, spectinomycin, azithromycin, and any other drugs in local use for gonorrhea treatment.

In cases of persistent gonococcal infection after treatment, clinicians should consider performing culture and antimicrobial susceptibility testing. In areas where fluoroquinolones are used for treating gonorrhea and small numbers of patients with QRNG are identified, health departments should notify and treat partners of patients with known QRNG to minimize the spread of resistance. Through their state and local health departments, clinicians and laboratorians should report treatment failures or resistant gonococcal isolates to CDC, telephone 404-639-8373; isolates may be submitted to CDC's *Neisseria* Reference Laboratory for confirmation testing, telephone 404-639-3470.

Acknowledgments

This report is based in part on assistance and data contributed by AM Vannier, MD, Southern California Kaiser Permanente Regional Reference Laboratories, Los Angeles, California; N O'Connor, Hawaii State Public Health Laboratory, Pearl City; P Effler, MD, Hawaii State Dept of Health; N DeAugustine, H Calvet, MD, M Lachica, City of Long Beach Dept of Health and Human Svcs, Long Beach; PR Kerndt, MD, Los Angeles County Dept of Health Svcs, Los Angeles; P Weismuller, DrPH, County of Orange Health Care Agency, Santa Ana; D Moore, PhD, P Hannah, Orange County Public Health Laboratory, Santa Ana; C Peter, PhD, D Kiefner, G Washabaugh, MPH, San Diego County Health and Human Svcs Agency, San Diego; JD Klausner, MD, L Fischer, MPA, V Zapitz, S Liska, DrPH, San Francisco Dept of Public Health, San Francisco; S Coulter, E Lopez, STD Control Br, California Dept of Health

Svcs. F Judson, MD, Denver Gonococcal Isolate Surveillance Project (GISP) Regional Laboratory, Denver Health, Colorado; KK Holmes, MD, J Hale, MS, W Whittington, PhD, K Winterscheid, MS, Seattle GISP Regional Laboratory, Univ of Washington, Seattle; JS Knapp, PhD, DL Trees, PhD, Div of AIDS, STD, and TB Laboratory Research, National Center for Infectious Diseases; AB Harvey, SM Conner, MPH, Div of STD Prevention, National Center for HIV, STD, and TB Prevention, CDC.

References

1. Fleming DT, Wasserheit JN. From epidemiological synergy to public health policy and practice: the contribution of other sexually transmitted diseases to sexual transmission of HIV infection. *Sex Transm Infect* 1999;75:3-17.
2. CDC. Sexually transmitted disease surveillance 2001. Atlanta, Georgia: U.S. Department of Health and Human Services, CDC, 2002.
3. CDC. Sexually transmitted disease surveillance 2001 supplement: Gonococcal Isolate Surveillance Project (GISP) Annual Report—2001. Atlanta, Georgia: U.S. Department of Health and Human Services, 2002.
4. Iverson CJ, Wang S, Ohye R, et al. Emergence of a possible endemic focus of ciprofloxacin-resistant *Neisseria gonorrhoeae* in Hawaii [Abstract]. In: Program and abstracts of the International Conference on Emerging Infectious Diseases. Atlanta, Georgia: U.S. Department of Health and Human Services, CDC, July 16-19, 2000.
5. WHO Western Pacific Gonococcal Antimicrobial Surveillance Programme. Surveillance of antibiotic resistance in *Neisseria gonorrhoeae* in the WHO Western Pacific Region, 2000. *Commun Dis Intell* 2001;25:274-7.
6. CDC. Sexually transmitted diseases treatment guidelines 2002. *MMWR* 2002;51(No. RR-6).
7. CDC. Discontinuation of cefixime tablets—United States. *MMWR* 2002;51:1052.
8. CDC. Fluoroquinolone-resistance in *Neisseria gonorrhoeae* in Hawaii, 1999, and decreased susceptibility to azithromycin in *N. gonorrhoeae*, Missouri, 1999. *MMWR* 2000;49:833-6.
9. CDC. Antibiotic-resistant strains of *Neisseria gonorrhoeae*: policy guidelines for detection, management, and control. *MMWR* 1987;36(No. S-5).
10. Roy K, Wang S, Meltzer M. Identifying the optimal diagnostic and treatment strategy for gonorrhea in a time of increasing ciprofloxacin-resistance [Abstract]. In: Program and abstracts of the 2002 Conference on Antimicrobial Resistance. Bethesda, Maryland: National Foundation for Infectious Diseases, June 27-29, 2002.

Multistate Outbreaks of *Salmonella* Serotype Poona Infections Associated with Eating Cantaloupe from Mexico — United States and Canada, 2000–2002

Three multistate outbreaks of *Salmonella* serotype Poona infections associated with eating cantaloupe imported from Mexico occurred in the spring of consecutive years during 2000–2002. In each outbreak, the isolates had indistinguishable pulsed-field gel electrophoresis (PFGE) patterns; the PFGE patterns observed in the 2000 and 2002 outbreaks were

indistinguishable, but the pattern from 2001 was unique among them. Outbreaks were identified first by the California Department of Health Services (2000 and 2001) and the Washington State Department of Health (2002) and involved residents of 12 states and Canada. This report describes the investigations, which led ultimately to an import alert on cantaloupes from Mexico. To limit the potential for cantaloupe contamination, the Food and Drug Administration (FDA) continues to work with the Mexican government on a food-safety program for the production, packing, and shipping of fresh cantaloupes.

April–June 2000 Outbreak

A total of 47 confirmed cases of *S. Poona* infections with indistinguishable PFGE patterns were identified from California (26), Washington (10), Nevada (five), New Mexico (three), Oregon (two), and Colorado (one), with illness onset occurring during April 14–June 2. The median age of ill persons was 7 years (range: 1–95 years); 28 (60%) patients were aged <10 years, and nine (19%) were aged >60 years. Twenty-four (51%) patients were male and nine (19%) were hospitalized.

A matched case-control study was conducted; 20 case-patients were matched by age category to 37 community controls. A case was defined as laboratory-confirmed infection with *S. Poona* of the outbreak PFGE pattern in a person with illness onset during April–June. By multivariable modeling, illness was associated only with eating cantaloupe (matched odds ratio [MOR]=6.7; 95% confidence interval [CI]=1.3–34.0), with 16 (80%) case-patients versus seven (19%) controls reporting eating cantaloupe. Cantaloupe was purchased either pre-cut or whole.

April–May 2001 Outbreak

In April, an initial cluster of *S. Poona* was identified in California. Isolates had a rare biochemical trait, the inability to produce hydrogen sulfide (H_2S), and PFGE patterns that were indistinguishable. A total of 50 cases of H_2S -negative *S. Poona* infections were identified in residents of California (28), Washington (eight), Nevada (seven), Arizona (six), and Oregon (one). Demographic and illness-history data from the 28 California patients indicated that illness onset occurred during April 6–May 28. The age distribution was bimodal; the 19 children had a median age of 3 years (range: 1–5 years) and the nine adults had a median age of 80 years (range: 39–91 years). Fifteen (54%) patients were female. Ten (36%) patients were bacteremic; one infant girl had *S. Poona* isolated from a urine specimen. Nine (33%) patients were

hospitalized, and two patients (a man aged 78 years and a woman aged 91 years) died with *Salmonella* septicemia.

A matched case-control study was conducted; 11 case-patients from California (seven), Nevada (two), Arizona (one), and Washington (one) were matched by age category to 19 community controls. Case-patients had laboratory-confirmed infections of the outbreak strain of H_2S -negative *S. Poona* and illness onset during the first 2 weeks of April. Illness was associated only with eating cantaloupe (MOR=7.4; 95% CI=1.0–178.0). Eight (80%) case-patients and six (33%) controls recalled eating cantaloupe. Cantaloupe was purchased either pre-cut or whole.

March–May 2002 Outbreak

A total of 58 cases with *S. Poona* isolates with indistinguishable PFGE patterns were identified in California (21), Washington (nine), Oregon (five), British Columbia (four), Colorado (three), Nevada (three), Manitoba (two), Missouri (two), Ontario (two), Saskatchewan (two), Texas (two), Arkansas (one), Minnesota (one), and Vermont (one). Illness onset occurred during March 30–May 31; the median age of patients was 6 years (range: 4 months–91 years); 32 (55%) were aged <10 years, and 11 (19%) were aged >60 years. A total of 31 (55%) were female. Ten patients were hospitalized.

A matched case-control study was conducted; 27 case-patients were matched by age category to 54 community controls. A case was defined as *S. Poona* infection with the outbreak PFGE pattern in a person aged ≥ 2 years with illness onset during March 15–May 3. The only exposure significantly associated with illness was eating cantaloupe; 20 (74%) case-patients recalled eating cantaloupe compared with 11 (20%) controls (MOR=15.5; 95% CI=3.3–125.0). Case-patients (50%) were more likely than controls (13%) to eat cantaloupe purchased whole (MOR=5.8; 95% CI=1.6–23.3) or to eat cantaloupe in a fruit salad or as a garnish (28% versus 5%) (MOR=6.5; 95% CI=1.2–63.0). No other factors were significantly associated with illness.

Traceback and Regulatory Action

FDA, in conjunction with state and provincial food regulatory agencies, conducted traceback investigations of cantaloupe purchased by patients in all three outbreaks. In each instance, point-of-sale sources of cantaloupe were traced back to shippers and then to farms in Mexico. In response to the 2000 and 2001 outbreaks, FDA conducted on-farm investigations in Mexico and concluded that measures were not in place to minimize microbial contamination in the growing, harvesting, packaging, and cooling of cantaloupe. Possible

sources of contamination include irrigation of fields with water contaminated with sewage, processing (cleaning and cooling) produce with *Salmonella*-contaminated water, poor hygienic practices of workers who harvest and process the cantaloupe, pests in packing facilities, and inadequate cleaning and sanitizing of equipment that comes in contact with cantaloupe. In association with the 2001 outbreak, FDA detained product imported by the shipper on May 31, and the shipper voluntarily recalled its imported Mexican cantaloupe. The shipper and the implicated farm in Mexico remain on detention. In association with the 2002 outbreak, the importer voluntarily recalled the implicated Mexican cantaloupe, and FDA placed the implicated farms on detention. On October 28, 2002, FDA issued an import alert on cantaloupe from Mexico that detains all products offered for entry at all U.S. ports.

Reported by: SM Anderson, MPH, Arizona Dept of Health Svcs. L Verchick, MS, Clark County Health Department, Las Vegas; R Sowadsky, MSPH, Nevada State Health Div. B Sun, DVM, R Civen, MD, JC Mohle-Boetani, MD, SB Werner, MD, M Starr, DVM, S Abbott, M Gutierrez, M Palumbo, PhD, J Farrar, PhD, California Dept of Health Svcs. P Shillam, Colorado Dept of Health. E Umland, MD, M Tanuz, M Sewell, DrPH, J Cato, New Mexico Dept of Health. W Keene, PhD, Oregon Dept of Human Svcs. M Goldoft, MD, J Hofmann, MD, J Kobayashi, MD, P Waller, MS, Washington State Dept of Health. Center for Food Safety and Applied Nutrition and the Office of Regulatory Affairs, Food and Drug Administration. C Braden, MD, Div of Bacterial and Mycotic Diseases, National Center for Infectious Diseases; G Djomand, MD, M Reller, MD, W Chege, MD, EIS officers, CDC.

Editorial Note: *Salmonella* infections have been linked to melons at least since 1990 when *Salmonella* serotype Chester traced to cantaloupe caused 245 illnesses in 30 states (1). The cantaloupe were imported from either Mexico or Guatemala. In 1991, an outbreak of cantaloupe-associated *S. Poona* infections caused 400 illnesses in 23 states (2). Illness was associated with eating pre-cut cantaloupe in fruit salads or from salad bars. Although industry sources identified the lower Rio Grande Valley in Texas as the probable source of the implicated cantaloupe, some might have come from Mexico. In response to this outbreak, FDA conducted a microbiologic survey that isolated a variety of *Salmonella* serotypes from approximately 1% of sampled imported cantaloupe and watermelon (2). In 1997, an outbreak of *Salmonella* serotype Saphra infections affected 25 persons in California. Illness was associated with cantaloupe imported from Mexico (3). After the 2000 and 2001 *S. Poona* outbreaks, FDA conducted farm investigations in Mexico, issued press releases to warn consumers, placed implicated farms on detention, and conducted sampling surveys of imported cantaloupe. The 1999 and 2000 FDA surveys of imported produce indicated

that 5% of cantaloupe sampled (eight of 151) was contaminated with *Salmonella* (4). A 2001 survey of imported produce indicates that of 29 cantaloupes from Mexico tested, none yielded *Salmonella*, *Shigella*, or *Escherichia coli* O157:H7 (FDA, unpublished data, 2001). The interpretation of the 2001 survey is limited by the small sample size.

S. Poona is a relatively rare serotype that is responsible for 1% of human *Salmonella* isolates reported in the United States in 2001; however, of the six cantaloupe-associated *Salmonella* outbreaks, four were attributed to infections with *S. Poona*. Typically, human infection with *S. Poona* is associated with reptile exposure (5,6). The three outbreaks attributed to *S. Poona*-contaminated cantaloupe traced to Mexican farms suggest the possibility of a unique natural reservoir in the Mexican farm environment, possibly from reptiles such as iguanas drawn to feed on melon crops that enter the packing sheds and contaminate the equipment. Subsequently, water used in the washing and cooling process might spread the contamination.

FDA provides information about the decontamination of melons to the retail industry, food-service establishments, and commercial processors of pre-cut melon (7,8). The use of sodium hypochlorite or other permitted antimicrobials in combination with brushing is recommended. The potential for microbial contamination also might be reduced by using only good-quality fruit that is free from open wounds or defects that might allow bacteria to contaminate the interior of the fruit (9). Additional research is needed to determine the effectiveness of consumer produce-washing practices. Consumers should be sure that fresh-cut melons are refrigerated or surrounded by ice; leftover cut melons should be discarded if left at room temperature for >2 hours. Additional information for consumers is available at <http://www.fda.gov/bbs/topics/answers/2002/ans01167.html>.

On October 28, 2002, in response to the three outbreaks during 2000–2002 and analytical results from the sampling of imported Mexican cantaloupe, FDA issued an import alert that detains all cantaloupe from Mexico offered for entry at all U.S. ports. FDA will continue to work with the Mexican government on a food-safety program for the production, packing, and shipping of fresh cantaloupe. The Mexican government is developing a certification program based on sound agricultural and manufacturing practices that would allow FDA to identify farms that have adopted and implemented such a food-safety program.

Acknowledgments

This report is based in part on assistance and data contributed by J Anderberg, Food Safety Program, Washington State Dept of Health. S Stenzel, K Smith, Minnesota Dept of Health. B Labus,

P Rowley, Clark County Health District, Las Vegas, Nevada. S Schoenfeld, Vermont Dept of Health. L Gaul, Texas Dept of Health. S Isaacs, A Ellis, Health Canada, Ottawa; M Fyfe, British Columbia Center for Disease Control, Vancouver; H Bangura, Saskatchewan Health, Regina, Canada. J Varma, J Painter, Div of Bacterial and Mycotic Diseases, National Center for Infectious Diseases, CDC.

References

1. Ries AA, Zaza S, Langkop C, et al. A multistate outbreak of *Salmonella* Chester linked to imported cantaloupe [Abstract]. In: Programs and abstracts of the 30th Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 1990.
2. CDC. Epidemiologic notes and reports: multistate outbreak of *Salmonella* Poona infections—United States and Canada, 1991. MMWR 1991;40:549–52.
3. Mohle-Boetani JC, Reporter R, Werner SB, et al. An outbreak of *Salmonella* serogroup Saphra due to cantaloupes from Mexico. J Infect Dis 1999;180:1361–4.
4. Food and Drug Administration. FDA survey of imported fresh produce: FY 1999 field assignment. Available at <http://www.cfsan.fda.gov/~dms/prodsur6.html>.
5. Reporter R, Bendana N, Sato H, et al. Rare serotypes of *Salmonella* associated with iguana exposure [abstract 1460]. In: Program and Abstracts of the 33rd Interscience Conference on Antimicrobial Agents and Chemotherapy. Washington, DC: American Society for Microbiology, 1999.
6. Woodward DL, Khakhria R, Johnson WM. Human salmonellosis associated with exotic pets. J Clin Microbiol 1997;35:2786–90.
7. National Archives and Records Administration. Code of Federal Regulations. Title 21, Part 173: secondary direct food additives permitted in food for human consumption. Revised April 2002. Available at <http://www.access.gpo.gov/cgi-bin/cfrassemble.cgi?title=200221>.
8. Food and Drug Administration. Produce safety at retail: safe handling practices for melons. Available at <http://www.cfsan.fda.gov/~ear/ret-mln.html>.
9. Food and Drug Administration. Guidance for industry: guide to minimize microbial food safety hazards for fresh fruits and vegetables, 1998. Available at <http://www.cfsan.fda.gov/~dms/prodguid.html>.

Enterovirus Surveillance — United States, 2000–2001

Enteroviruses are common viruses associated with diverse clinical manifestations ranging from mild febrile illness to severe and potentially fatal syndromes including aseptic meningitis, encephalitis, neonatal systemic enteroviral disease, and paralytic poliomyelitis (1). A total of 64 enterovirus serotypes are recognized, including 61 nonpolio enteroviruses (2)*. Individual serotypes have

different temporal patterns of circulation and often are associated with different clinical manifestations (1,4). This report describes temporal trends in reported enterovirus infections in the United States during 2000–2001, including widespread activity of two serotypes (echoviruses 13 and 18) that previously were detected rarely. Monitoring of circulating enterovirus serotypes helped identify these two agents as the primary causes of aseptic meningitis outbreaks in the United States in 2001. Further improvements in timeliness of reporting and geographic representation of the system are needed to allow more complete surveillance for enteroviruses.

Other than paralytic polio, diseases associated with enterovirus infections, including aseptic meningitis, are not nationally notifiable in the United States. The National Enterovirus Surveillance System (NESS) collects information on enterovirus serotypes and monitors temporal and geographic trends to help public health officials recognize and control outbreaks of enteroviral disease. Enterovirus detections from human specimens that are submitted for testing to participating laboratories are reported voluntarily to NESS along with basic demographic information, specimen type, and date of collection.

The number of laboratories participating in NESS increased from eight in 1999 to 21 in 2000 and 25 in 2001. During 2000–2001, a total of 27 laboratories participated in NESS, including 24 state public health laboratories, two private laboratories, and CDC's Enterovirus Laboratory, which receive specimens from multiple states. Enterovirus detections were reported in 36 states in 2000 and in 30 states in 2001. During 2000–2001, a total of 40 states reported enterovirus detections; of 2,319 reports, 1,925 (83.0%) were submitted by public health laboratories, 318 (13.7%) by private laboratories, and 76 (3.3%) by CDC. Of the 27 laboratories, three used genomic sequencing for enterovirus typing, and 24 used traditional antigenic typing methods. Serotypes were identified for 1,863 (80.3%) reports (Table 1). Consistent with the trend observed throughout the 1990s, the proportion of reported enteroviruses with unknown serotype increased from 13.1% during 1997–1999 to 19.7% during 2000–2001. Because the high proportion of unknown serotypes could lead

TABLE 1. Number and percentage of enterovirus detection reports, by serotype identification status — National Enterovirus Surveillance System, United States, 2000–2001

| Serotype status | 2000 | | Years | | Total | |
|------------------------|------------|---------|--------------|---------|--------------|---------|
| | No. | (%) | No. | (%) | No. | (%) |
| With known serotypes | 578 | (78.6%) | 1,285 | (81.1%) | 1,863 | (80.3%) |
| With unknown serotypes | 157 | (21.4%) | 299 | (18.9%) | 456 | (19.7%) |
| Total | 735 | | 1,584 | | 2,319 | |

* Echoviruses 22 and 23 have been reclassified recently as human parechoviruses 1 and 2, respectively, members of genus *Parechovirus*, which is related to (but distinct from) the genus *Enterovirus*. Epidemiologic and clinical features of these viruses are similar to those of the enteroviruses (3), and detection of these viruses continues to be reported to the National Enterovirus Surveillance System.

to underestimating the number of individual enteroviruses, reports with unknown serotypes were excluded from the analysis of serotype distribution.

During 2000–2001, echovirus 18 and echovirus 13 were the predominant serotypes, accounting for 22.0% and 20.8% respectively of the reports with an identified serotype, followed by coxsackievirus B5 (11.9%), coxsackievirus B2 (6.3%), and echovirus 6 (6.1%) (Table 2). The serotype detected most commonly for 2000 was coxsackievirus B5 (34.4%). The predominant enteroviruses in 2001 were echoviruses 18 and 13, which accounted for 30.8% and 29.3%, respectively; echovirus 13 was reported in 24 states and echovirus 18 in 19 states. Illinois, Michigan, Tennessee, and Wisconsin reported the most echovirus 13 detections, and Illinois, Minnesota, New York, and Texas reported the most echovirus 18 detections. One vaccine-related type 3 poliovirus was reported in 2000.

The most common source for enterovirus detection was cerebrospinal fluid (980 [51.2%] of 1,915 reports indicating the source specimen), followed by stool or rectal swab (338 [17.7%]). Children aged <1 year accounted for 859 (44.0%) of 1,951 reports for which age data were available.

Reported by: State virology laboratory directors, Diagnostic Virology Laboratory, Associated Regional and Univ Pathologists Laboratories, Salt Lake City, Utah, Diagnostic Virology Laboratory, Texas Children's Hospital, Houston, Texas, S Oberste, PhD, A LaMonte, MPH, N Khetsuriani, MD, Div of Viral and Rickettsial Diseases, National Center for Infectious Diseases; V Hsu, MD, J Mullins, DVM, EIS officers, CDC.

TABLE 2. Distribution of the 15 most commonly reported nonpolio enterovirus serotypes, by rank — National Enterovirus Surveillance System, United States, 2000–2001

| Rank | 2000 (n=577) | | 2001 (n=1,285) | | 2000–2001 (n=1,862) | |
|---------------|-----------------|-------------|-------------------|-------------|------------------------|-------------|
| | Serotype | % | Serotype | % | Serotype | % |
| 1 | coxsackie B5 | 34.4 | echo 18 | 30.8 | echo 18 | 22.0 |
| 2 | echo 6 | 8.8 | echo 13 | 29.3 | echo 13 | 20.8 |
| 3 | coxsackie A9 | 8.7 | coxsackie B2 | 7.6 | coxsackie B5 | 11.9 |
| 4 | coxsackie B4 | 8.3 | echo 6 | 4.8 | coxsackie B2 | 6.3 |
| 5 | echo 11 | 6.9 | echo 4 | 4.1 | echo 6 | 6.1 |
| 6 | echo 9 | 6.2 | echo 11 | 3.4 | echo 11 | 4.5 |
| 7 | coxsackie B2 | 3.5 | coxsackie B3 | 3.0 | coxsackie A9 | 4.0 |
| 8 | echo 25 | 2.6 | coxsackie B1 | 2.7 | echo 9 | 3.3 |
| 9 | echo 18 | 2.3 | echo 9 | 2.0 | coxsackie B4 | 3.2 |
| 10 | enterovirus 71 | 2.1 | coxsackie A9 | 2.0 | echo 4 | 3.1 |
| 11 | echo 16 | 1.9 | coxsackie B5 | 1.7 | coxsackie B3 | 2.4 |
| 12 | echo 30 | 1.9 | echo 30 | 1.7 | coxsackie B1 | 2.0 |
| 13 | echo 13 | 1.7 | coxsackie B4 | 0.9 | echo 30 | 1.8 |
| 14 | echo 21 | 1.6 | echo 25 | 0.6 | echo 25 | 1.2 |
| 15 | parecho 1* | 1.4 | enterovirus 71 | 0.6 | enterovirus 71 | 1.1 |
| Total† | | 92.2 | | 95.3 | | 93.5 |

* Formerly echo 22.

† Totals might be slightly different from sums of percentages because of rounding. For all other serotypes, percentages were 7.8% in 2000, 4.7% in 2001, and 6.5% during 2000–2001.

Editorial Note: Serotype-based enterovirus surveillance in the United States has five objectives. First, NESS data help to determine long-term patterns of circulation for individual serotypes (4). Second, the data are used to associate trends in enteroviral diseases with circulating serotypes such as viral meningitis-associated hospitalizations during periods of high activity of echoviruses 9 and 30 and lower numbers of these cases for years when group B coxsackieviruses predominate (CDC, unpublished data, 1988–1999). Third, the data are used to guide outbreak investigations. Fourth, because different serotypes have differential sensitivity to at least one candidate antienterovirus drug (5), information on circulating serotypes helps guide the development of new diagnostic tests and therapies. Finally, NESS monitors circulation of poliovirus strains to supplement poliovirus surveillance; this aspect of enterovirus surveillance will remain important until successful global poliovirus eradication ends the need for polio surveillance.

The findings in this report are consistent with previous observations on temporal variability of predominant serotypes. Of the 15 serotypes reported most commonly during 2000–2001, seven (coxsackieviruses A9, B2, and B4 and echoviruses 6, 9, 11, and 30) have appeared consistently among the 15 most common serotypes each year during 1993–1999 (6,7). Two enteroviruses (echovirus 18 and echovirus 13) that previously were rarely reported emerged as the predominant serotypes in 2001. Timely identification of increased activity by these serotypes helped guide investigation of outbreaks of aseptic meningitis reported to CDC in 2001 from six states (Alaska, Louisiana, Maryland, Mississippi, Montana, and Tennessee); all of these outbreaks were linked subsequently to one or both of these serotypes (8). For echovirus 13, this was the first report of widespread circulation in the United States and probably reflected the worldwide activity of this serotype that has been observed since 2000 (8).

Multiple factors might explain the increase in unknown serotypes, including limited availability of the appropriate typing antisera, high cost of reagents, and labor intensity of testing. The use of enterovirus typing based on genomic sequencing (9) could increase the proportion of identified serotypes, including those for which reagents are not readily available. The virtual absence of vaccine-related poliovirus isolates after 2000 is associated with the discontinuation of the use of oral polio vaccine in the United States beginning in 2000 (10).

The findings in this report are subject to at least two limitations. First, because of the voluntary and passive nature of NESS, the small number of reports for some states, and the lack of reporting or testing by others, these results might not be fully representative of the entire United States. Second, because many serotypes were not identified, the number of individual enteroviruses might be underestimated.

As laboratory participation in NESS increases, the data will become more representative geographically. More timely reporting from laboratories would allow NESS to provide frequent feedback in the form of an online enterovirus surveillance summary, which would increase the public health utility of this surveillance system.

References

1. Pallansch MA, Roos RP. Enteroviruses: polioviruses, coxsackieviruses, echoviruses, and newer enteroviruses. In: Knipe DM, Howley PM, Griffin DE, et al. eds. *Fields Virology*. 4th ed. Philadelphia, Pennsylvania: Lippincott-Raven Publishers, 2001.
2. Institute for Animal Health. The classification of the *picornaviridae*. Compton, United Kingdom: Institute for Animal Health, 2002. Available at <http://www.iah.bbsrc.ac.uk/virus/picornaviridae/picornastudygroup/ictv-7a.htm>.
3. Stanway G, Joki-Korpela P, Hyypia T. Human parechoviruses—biology and clinical significance. *Rev Med Virol* 2000;10:57–69.
4. Strikas RA, Anderson L, Parker RA. Temporal and geographic patterns of isolates of nonpolio enteroviruses in the United States, 1970–1983. *J Infect Dis* 1986;153:346–51.
5. Pevar DC, Tull TM, Seipel ME, Groarke JM. Activity of pleconaril against enteroviruses. *Antimicrob Agents Chemother* 1999;43:2109–15.
6. CDC. Nonpolio enterovirus surveillance—United States, 1993–1996. *MMWR* 1997;46:748–50.
7. CDC. Enterovirus surveillance—United States, 1997–1999. *MMWR* 1999;49:913–6.
8. Khetsuriani N, Oberste S, Mullins J, et al. Changes in circulating enterovirus serotypes in the United States—United States, 2001 [Abstract]. Presented at the International Conference on Emerging Infectious Diseases, Atlanta, Georgia: US Department of Health and Human Services, CDC, March 24–27, 2002.
9. Oberste MS, Maher K, Kilpatrick DR, Pallansch MA. Molecular evolution of the human enteroviruses: correlation of serotype with VP1 sequence and application to picornavirus classification. *J Virol* 1999;73:1941–8.
10. CDC. Poliomyelitis prevention in the United States: updated recommendations of the Advisory Committee on Immunization Practices (ACIP). *MMWR* 2000;49(No. RR-5).

West Nile Virus Activity — United States, November 14–20, 2002, and Missouri, January 1– November 9, 2002


This report summarizes West Nile virus (WNV) surveillance data reported to CDC through ArboNET and by states and other jurisdictions as of 9 a.m. Mountain Standard Time, November 20, 2002.

United States

During November 14–20, a total of 111 laboratory-positive human cases of WNV-associated illness were reported from Indiana (n=37), Illinois (n=23), Texas (n=17), Nebraska (n=nine), Ohio (n=six), Georgia (n=five), Arkansas (n=four), Florida (n=four), Minnesota (n=three), Maryland (n=one), Massachusetts (n=one), and Tennessee (n=one). During the same period, WNV infections were reported in 90 dead crows and 330 other dead birds. A total of 411 veterinary cases and 25 WNV-positive mosquito pools were reported.

During 2002, a total of 3,698 human cases with laboratory evidence of recent WNV infection have been reported from Illinois (n=761), Michigan (n=504), Ohio (n=419), Louisiana (n=323), Indiana (n=284), Mississippi (n=182), Missouri (n=169), Texas (n=165), Nebraska (n=124), New York (n=78), Kentucky (n=67), Pennsylvania (n=59), Tennessee (n=55), Iowa (n=48), Alabama (n=46), Minnesota (n=45), Wisconsin (n=42), South Dakota (n=37), Georgia (n=35), the District of Columbia (n=34), Maryland (n=29), Virginia (n=27), Arkansas (n=25), Massachusetts (n=23), Florida (n=22), Connecticut (n=17), North Dakota (n=17), Oklahoma (n=16), Colorado (n=12), New Jersey (n=12), Kansas (n=nine), West Virginia (n=three), North Carolina (n=two), California (n=one), Delaware (n=one), Montana (n=one), Rhode Island (n=one), South Carolina (n=one), Vermont (n=one), and Wyoming (n=one) (Figure 1). Among the 3,287 patients for whom data were available, the median age was 55 years (range: 1.5 months–99 years); 1,755 (54%) were male, and the dates of illness onset ranged from June 10 to

FIGURE 1. Areas reporting West Nile virus (WNV) activity — United States, 2002*



* As of 9 a.m. Mountain Standard Time, November 20, 2002.

† California has reported human WNV activity only.

November 4. A total of 198 human deaths have been reported. The median age of decedents was 78 years (range: 24–99 years); 120 (61%) deaths were among men. In addition, 7,612 dead crows and 6,060 other dead birds with WNV infection were reported from 42 states and the District of Columbia; 8,723 WNV infections in mammals (8,710 equines, three canines, and 10 other species) have been reported from 38 states (Alabama, Arkansas, Colorado, Connecticut, Delaware, Florida, Georgia, Idaho, Illinois, Indiana, Iowa, Kansas, Kentucky, Louisiana, Maryland, Massachusetts, Minnesota, Mississippi, Missouri, Montana, Nebraska, New Jersey, New Mexico, New York, North Carolina, North Dakota, Ohio, Oklahoma, Pennsylvania, South Carolina, South Dakota, Tennessee, Texas, Vermont, Virginia, Washington, Wisconsin, and Wyoming). During 2002, WNV seroconversions have been reported in 366 sentinel chicken flocks from Florida, Iowa, Nebraska, North Carolina, Pennsylvania, Texas, and New York City; 4,931 WNV-positive mosquito pools have been reported from 27 states (Alabama, Arkansas, Connecticut, Delaware, Georgia, Illinois, Indiana, Iowa, Kansas, Kentucky, Maryland, Massachusetts, Mississippi, Missouri, Nebraska, New Hampshire, New Jersey, New York, North Carolina, Ohio, Pennsylvania, Rhode Island, South Carolina, South Dakota, Texas, Vermont, and Virginia), New York City, and the District of Columbia.

Missouri

During January 1–November 9, 2002, the Missouri Department of Health and Senior Services (MDHSS) identified 168 persons with laboratory evidence of West Nile virus (WNV) infection; 34 cases were confirmed by CDC on the basis of the plaque-reduction neutralization test, and 134 cases were classified as probable pending CDC confirmation. Five cases were fatal; all five patients had encephalitis.

The 168 patients had a median age of 53.2 years (range: 4–92 years); 91 patients (53%) were male. Median age of decedents was 75.6 years (range: 61–88 years). Dates of illness onset ranged from July 31 to October 12.

Of Missouri's 114 reporting local public health agencies (LPHAs), 109 (96%) have reported WNV-positive animal tests, representing nearly all geographic regions and human population centers in Missouri (Figure 2). Human WNV cases have occurred among persons from 32 (28%) of LPHA districts; 126 (75%) patients resided in metropolitan St. Louis districts. The metropolitan Kansas City area has recorded eight cases; no human WNV cases have been reported from the Springfield area (Greene County).

The elevated WNV incidence rate in the metropolitan St. Louis area, especially within St. Louis city limits (16.1

FIGURE 2. Number of West Nile virus cases in humans*, by county — Missouri, January 1–November 9, 2002



* n=168.

cases per 100,000 population), is under investigation. Of 134 Missouri WNV cases for which case investigations are completed, 86 (64%) patients recalled being bitten by a mosquito 3 weeks before onset of illness. In addition, 86 (64%) patients reported not using repellent during that time. Patients spent an average of 4.4 hours outdoors per day for leisure and work. Morning (47%) was the time that patients most frequently reported being active outdoors for leisure and work. A total of 58 (43%) patients reported having standing water around their home during the 3 weeks preceding onset of illness. The top four reported sources of standing water were birdbaths, flowerpots, pet watering containers, and tires.

MDHSS provides frequent updates on human cases of WNV and activity by county for dead birds and infected horses on its website (<http://www.dhss.state.mo.us/westnilevirus/positives.pdf>). Results on specimens submitted for laboratory testing are provided to patient providers, submitting laboratories, and local health departments. MDHSS issues press releases updating case numbers, reminding the public of personal protection measures, and advising elimination of sources of standing water around residences. MDHSS produced a statewide prevention campaign based on the slogan "Don't Let It Bug You," which included posters and brochures distributed to all 114 LPHAs, direct-mail appeals to St. Louis area clergy for increased community awareness of mosquito habitats, television and radio public service announcements, and a toll-free hotline for citizens' questions. Mosquito-control program recommendations, developed by MDHSS

are available at <http://www.dhss.state.mo.us>. The decision to initiate a control program has been left to local municipalities.

Additional information about WNV activity is available at <http://www.cdc.gov/ncidod/dvbid/westnile/index.htm> and http://www.cindi.usgs.gov/hazard/event/west_nile/west_nile.html.

Notice to Readers

Approval of a New Rapid Test for HIV Antibody

On November 7, 2002, the Food and Drug Administration announced approval of the OraQuick Rapid HIV-1 Antibody Test (OraSure Technologies, Inc., Bethlehem, Pennsylvania) for use by trained personnel as a point-of-care test to aid in the diagnosis of infection with human immunodeficiency virus type 1 (HIV-1). OraQuick is a simple, rapid test that can detect antibodies to HIV in fingerstick whole blood specimens and provide results in ≤ 20 minutes. The test has been categorized as moderate complexity under the Clinical Laboratory Improvement Amendments of 1988 (CLIA). A second FDA-approved moderate-complexity rapid HIV test, Single Use Diagnostic System for HIV-1 (Abbott-Murex Inc., Norcross, Georgia), remains available in the United States for use with serum or plasma specimens.

Use of a rapid test that allows same-day results can substantially increase the number of persons who receive their test results, which improves the delivery of counseling and treatment services (1). On the basis of data submitted by the manufacturer for test approval, the sensitivity* of OraQuick in the clinical studies performed was 99.6% (95% confidence interval [CI]=98.5%–99.9%), and specificity was 100% (95% CI=99.7%–100%), comparable to those of FDA-approved enzyme immunoassays in widespread use. Because HIV prevalence is low in most U.S. testing settings, the negative predictive value† of screening with a single rapid test is high. Therefore, a negative rapid HIV test does not require further testing, and negative results with counseling can be provided at the initial visit. Retesting is recommended for those persons with a recent (within 3 months) history of known or possible exposure to HIV because there might have been insufficient time for detectable antibodies to develop (2). As with any HIV screening test, all reactive (preliminary positive) rapid test results should be confirmed by supplemental testing by either a Western blot or immunofluorescence assay

(3). The confirmatory tests can be performed on serum specimens obtained by phlebotomy, dried blot spots obtained on filter paper, or oral fluid specimens collected with the OraSure collection device.

Persons whose rapid-test results are reactive should be counseled about their likelihood of being infected with HIV and precautions to prevent HIV transmission, but they should return for definitive test results before medical referrals or partner counseling is initiated (3). A simple message to convey this information could be a statement that "Your preliminary test result was positive, but we won't know for sure if you are HIV-infected until we get the results from your confirmatory test. In the meantime, you should take precautions to avoid possibly transmitting the virus."

The Public Health Service recommends that rapid HIV tests should be used and preliminary positive test results provided when tested persons might benefit (1). Decisions about whether to use rapid tests should be based on considerations of return rates for standard test results and urgency of the need for test results (i.e., when necessary to make decisions about postexposure or perinatal prophylaxis) (1,4,5). The use of rapid tests will facilitate the acceptance of HIV testing and improve receipt of results in other health-care settings in which HIV testing is recommended, such as hospitals and acute-care clinics, where persons who are unaware of their HIV status might seek health-care services (6). Additional information and guidance on the use of rapid HIV tests are available from CDC at <http://www.cdc.gov/hiv/testing.htm>.

Sites wanting to perform this new HIV-1 rapid test that are not already certified to perform moderate-complexity laboratory tests under CLIA must enroll in the CLIA program, administered by the Centers for Medicare and Medicaid Services. The application and state agency contact information are available at <http://www.cms.hhs.gov/clia>. Information about enrollment and the requirements for moderate-complexity testing are available at <http://www.phppo.cdc.gov/clia/default.asp>.

CLIA moderate-complexity requirements provide minimum standards for personnel, quality control, proficiency testing, and quality assurance. In addition, some states have specific requirements that might apply to laboratory testing in general or to HIV testing specifically.

References

1. CDC. Update: HIV counseling and testing using rapid tests—United States, 1995. *MMWR* 1998;47:211–5.
2. CDC. Interpretation and use of the Western blot assay for serodiagnosis of human immunodeficiency virus type 1 infections. *MMWR* 1989;38(suppl 7):S4–S6.
3. CDC. Revised guidelines for HIV counseling, testing, and referral. *MMWR* 2001;50(No. RR-19).

* Sensitivity is the probability that the test result will be reactive if the specimen is a true positive; specificity is the probability that the test result will be nonreactive if the specimen is a true negative.

† The predictive value of a screening test is the probability that the test accurately predicts the true infection status of the person tested.

4. CDC. Updated U.S. Public Health Service guidelines for the management of occupational exposures to HBV, HCV, and HIV and recommendations for post-exposure prophylaxis. MMWR 2001;50(No. RR-11).
5. CDC. Revised recommendations for HIV screening of pregnant women. MMWR 2001;50(No. RR-19).
6. CDC. Recommendations for HIV testing services for inpatients and outpatients in acute-care hospital settings. MMWR 1993;42(No. RR-2).

Notice to Readers

Discontinuation of Cefixime Tablets — United States

In July 2002, Wyeth Pharmaceuticals (Collegeville, Pennsylvania) discontinued manufacturing cefixime (Suprax®) in the United States. In October 2002, the company ceased marketing cefixime tablets (200 mg and 400 mg) because of depletion of company inventory. Wyeth's patent for cefixime expired on November 10, 2002. No other pharmaceutical company manufactures or sells cefixime tablets in the United States. Wyeth will continue to sell cefixime suspension (100 mg/5 ml) until March 31, 2003, or until company inventory is depleted, whichever is sooner.

Cefixime is the only CDC-recommended oral antimicrobial agent to which *Neisseria gonorrhoeae* has not developed significant resistance (1). Uncomplicated *N. gonorrhoeae* infections may be treated with single-dose regimens of cefixime 400 mg orally, ceftriaxone 125 mg intramuscularly, or an oral fluoroquinolone (ciprofloxacin 500 mg, levofloxacin 250 mg, or ofloxacin 400 mg). However, fluoroquinolones should not be used for treatment of gonorrhea if the infection was acquired in Asia, the Pacific Islands (including Hawaii), or California because the prevalence of fluoroquinolone-resistant *N. gonorrhoeae* is high in those areas (1,2).

In the absence of cefixime, the primary recommended treatment option for gonorrhea in Hawaii and California is ceftriaxone. Also, in the absence of cefixime, ceftriaxone is the only CDC-recommended gonorrhea treatment option for young children and pregnant women throughout the United States. Fluoroquinolones can continue to be used for treating gonorrhea in areas of the United States with low prevalence of fluoroquinolone-resistant *N. gonorrhoeae*, but antimicrobial susceptibility monitoring should routinely be performed (2). Other oral antimicrobial agents, such as cefpodoxime,

cefuroxime axetil, and azithromycin, are not recommended by CDC for the treatment of gonorrhea. Additional information on the use of oral antimicrobials in treating *N. gonorrhoeae* infections will be available from CDC at <http://www.cdc.gov/std>.

References

1. CDC. Guidelines for treatment of sexually transmitted diseases 2002. MMWR 2002;51(No. RR-6).
2. CDC. Increases in fluoroquinolone-resistant *Neisseria gonorrhoeae*—Hawaii and California, 2001. MMWR 2002;51:1041–4.

Erratum: Vol. 51, No. 45

In the report "Influenza Outbreak—Madagascar, July–August, 2002," in Figure 2 on page 1017, the number of districts with laboratory-confirmed influenza was incorrectly indicated. Because a limited number of patients in a few districts were tested for influenza, the map has been changed to depict only districts with reported influenza-like illness cases.

FIGURE 2. Districts reporting cases of influenza-like illness (ILI) — Madagascar, July–August 2002





FIGURE I. Selected notifiable disease reports, United States, comparison of provisional 4-week totals ending November 16, 2002, with historical data

* No rubella cases were reported for the current 4-week period yielding a ratio for week 46 of zero (0).

[†] Ratio of current 4-week total to mean of 15 4-week totals (from previous, comparable, and subsequent 4-week periods for the past 5 years). The point where the hatched area begins is based on the mean and two standard deviations of these 4-week totals.

TABLE I. Summary of provisional cases of selected notifiable diseases, United States, cumulative, week ending November 16, 2002 (46th Week)*

| | Cum. 2002 | Cum. 2001 | | Cum. 2002 | Cum. 2001 |
|---|--------------|--------------|---|--------------|--------------|
| Anthrax | 2 | 21 | Encephalitis: West Nile [†] | 1,339 | 52 |
| Botulism: foodborne | 12 | 33 | Hansen disease (leprosy) [†] | 59 | 60 |
| infant | 48 | 86 | Hantavirus pulmonary syndrome [†] | 13 | 7 |
| other (wound & unspecified) | 26 | 16 | Hemolytic uremic syndrome, postdiarrheal [†] | 172 | 163 |
| Brucellosis [†] | 68 | 115 | HIV infection, pediatric ^{‡§} | 116 | 172 |
| Chancroid | 61 | 31 | Plague | - | 2 |
| Cholera | 5 | 4 | Poliomyelitis, paralytic | - | - |
| Cyclosporiasis [†] | 158 | 141 | Psittacosis [†] | 17 | 20 |
| Diphtheria | 1 | 2 | Q fever [†] | 43 | 23 |
| Ehrlichiosis: human granulocytic (HGE) [†] | 314 | 203 | Rabies, human | 2 | 1 |
| human monocytic (HME) [†] | 158 | 103 | Streptococcal toxic-shock syndrome [†] | 72 | 68 |
| other and unspecified | 9 | 5 | Tetanus | 20 | 27 |
| Encephalitis: California serogroup viral [†] | 117 | 112 | Toxic-shock syndrome | 100 | 105 |
| eastern equine [†] | 3 | 8 | Trichinosis | 12 | 21 |
| Powassan [†] | - | - | Tularemia [†] | 56 | 124 |
| St. Louis [†] | 8 | 76 | Yellow fever | 1 | - |
| western equine [†] | 2 | - | | | |

-: No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

[†] Not notifiable in all states.

[§] Updated monthly from reports to the Division of HIV/AIDS Prevention — Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention (NCHSTP). Last update October 31, 2002.

TABLE II. Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | AIDS | | Chlamydia [†] | | Cryptosporidiosis | | Escherichia coli, Enterohemorrhagic | | | |
|----------------|------------------------|-----------|------------------------|-----------|-------------------|-----------|-------------------------------------|-----------|--|-----------|
| | | | | | | | O157:H7 | | Shiga Toxin Positive, Serogroup non-O157 | |
| | Cum. 2002 [‡] | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| UNITED STATES | 24,713 | 34,080 | 681,846 | 684,617 | 2,555 | 3,458 | 3,220 | 2,884 | 150 | 142 |
| NEW ENGLAND | 1,011 | 1,268 | 23,786 | 21,449 | 167 | 137 | 252 | 230 | 32 | 38 |
| Maine | 23 | 40 | 1,466 | 1,187 | 11 | 18 | 37 | 26 | 5 | 1 |
| N.H. | 20 | 31 | 1,398 | 1,228 | 29 | 15 | 32 | 31 | - | 3 |
| Vt. | 8 | 13 | 835 | 544 | 31 | 32 | 12 | 13 | 1 | 1 |
| Mass. | 519 | 654 | 9,650 | 9,100 | 60 | 52 | 113 | 112 | 9 | 10 |
| R.I. | 71 | 84 | 2,443 | 2,602 | 20 | 4 | 14 | 13 | - | 1 |
| Conn. | 370 | 446 | 7,994 | 6,788 | 16 | 16 | 44 | 35 | 17 | 22 |
| MID. ATLANTIC | 5,619 | 8,977 | 76,678 | 74,744 | 312 | 316 | 220 | 218 | - | - |
| Upstate N.Y. | 404 | 1,168 | 14,968 | 12,534 | 126 | 94 | 160 | 139 | - | - |
| N.Y. City | 3,210 | 4,773 | 24,569 | 26,379 | 120 | 114 | 12 | 16 | - | - |
| N.J. | 925 | 1,509 | 10,764 | 12,416 | 10 | 19 | 48 | 63 | - | - |
| Pa. | 1,080 | 1,527 | 26,377 | 23,415 | 56 | 89 | N | N | - | - |
| E.N. CENTRAL | 2,494 | 2,499 | 119,570 | 127,395 | 817 | 1,509 | 778 | 749 | 19 | 11 |
| Ohio | 453 | 476 | 28,881 | 33,798 | 119 | 164 | 146 | 204 | 15 | 9 |
| Ind. | 347 | 306 | 15,168 | 13,712 | 51 | 78 | 68 | 79 | 1 | - |
| Ill. | 1,170 | 1,110 | 32,255 | 38,393 | 85 | 478 | 164 | 162 | - | - |
| Mich. | 398 | 457 | 28,812 | 26,784 | 112 | 173 | 131 | 90 | 3 | 2 |
| Wis. | 126 | 150 | 14,454 | 14,708 | 450 | 616 | 269 | 214 | - | - |
| W.N. CENTRAL | 421 | 718 | 37,450 | 34,775 | 386 | 497 | 483 | 464 | 37 | 38 |
| Minn. | 90 | 118 | 8,408 | 7,270 | 201 | 168 | 155 | 188 | 32 | 29 |
| Iowa | 54 | 80 | 4,761 | 4,451 | 42 | 80 | 115 | 76 | - | - |
| Mo. | 189 | 337 | 13,525 | 12,440 | 32 | 48 | 69 | 59 | N | N |
| N. Dak. | 1 | 2 | 801 | 901 | 20 | 13 | 17 | 19 | - | 2 |
| S. Dak. | 3 | 23 | 1,884 | 1,581 | 28 | 7 | 39 | 41 | 2 | 6 |
| Nebr. | 43 | 72 | 2,456 | 2,859 | 47 | 178 | 54 | 59 | 3 | 1 |
| Kans. | 41 | 86 | 5,615 | 5,273 | 16 | 3 | 34 | 22 | - | - |
| S. ATLANTIC | 7,537 | 10,268 | 131,239 | 131,527 | 316 | 344 | 347 | 224 | 36 | 33 |
| Del. | 131 | 217 | 2,363 | 2,488 | 3 | 6 | 8 | 4 | - | 1 |
| Md. | 1,066 | 1,517 | 14,542 | 13,547 | 21 | 36 | 25 | 28 | - | - |
| D.C. | 371 | 733 | 3,036 | 2,865 | 4 | 11 | - | - | - | - |
| Va. | 538 | 843 | 14,698 | 16,177 | 21 | 24 | 57 | 48 | 9 | 5 |
| W. Va. | 58 | 71 | 2,081 | 2,102 | 2 | 2 | 9 | 10 | - | - |
| N.C. | 555 | 778 | 22,110 | 19,279 | 32 | 27 | 130 | 46 | - | - |
| S.C. | 547 | 612 | 10,607 | 13,470 | 6 | 7 | 5 | 16 | - | - |
| Ga. | 1,160 | 1,232 | 26,326 | 28,506 | 133 | 150 | 54 | 43 | 10 | 9 |
| Fla. | 3,111 | 4,265 | 35,476 | 33,093 | 94 | 81 | 59 | 29 | 17 | 18 |
| E.S. CENTRAL | 1,128 | 1,532 | 41,595 | 44,064 | 109 | 48 | 99 | 127 | - | - |
| Ky. | 173 | 299 | 7,818 | 8,040 | 8 | 5 | 30 | 63 | - | - |
| Tenn. | 483 | 488 | 14,092 | 12,804 | 52 | 13 | 44 | 37 | - | - |
| Ala. | 197 | 378 | 11,034 | 12,536 | 42 | 16 | 18 | 16 | - | - |
| Miss. | 275 | 367 | 8,651 | 10,684 | 7 | 14 | 7 | 11 | - | - |
| W.S. CENTRAL | 2,696 | 3,435 | 94,418 | 94,900 | 35 | 123 | 64 | 183 | - | - |
| Ark. | 163 | 176 | 6,381 | 6,575 | 8 | 8 | 10 | 15 | - | - |
| La. | 693 | 699 | 16,946 | 16,254 | 5 | 7 | 2 | 7 | - | - |
| Okla. | 133 | 204 | 9,496 | 9,458 | 16 | 14 | 22 | 31 | - | - |
| Tex. | 1,707 | 2,356 | 61,595 | 62,613 | 6 | 94 | 30 | 130 | - | - |
| MOUNTAIN | 790 | 1,175 | 41,253 | 40,729 | 149 | 225 | 335 | 267 | 18 | 16 |
| Mont. | 8 | 15 | 1,976 | 1,677 | 5 | 37 | 29 | 20 | - | - |
| Idaho | 18 | 19 | 2,228 | 1,761 | 29 | 21 | 48 | 64 | 8 | 3 |
| Wyo. | 6 | 3 | 823 | 735 | 9 | 7 | 14 | 9 | 2 | 2 |
| Colo. | 157 | 262 | 12,003 | 11,737 | 53 | 39 | 87 | 87 | 4 | 6 |
| N. Mex. | 53 | 133 | 5,739 | 5,398 | 18 | 28 | 12 | 14 | 3 | 5 |
| Ariz. | 327 | 446 | 12,947 | 12,848 | 17 | 7 | 34 | 27 | 1 | - |
| Utah | 43 | 98 | 2,354 | 2,245 | 14 | 80 | 83 | 31 | - | - |
| Nev. | 178 | 199 | 3,183 | 4,328 | 4 | 6 | 28 | 15 | - | - |
| PACIFIC | 3,017 | 4,208 | 115,857 | 115,034 | 264 | 259 | 642 | 422 | 8 | 6 |
| Wash. | 302 | 427 | 13,010 | 12,010 | 43 | U | 138 | 115 | - | - |
| Oreg. | 216 | 177 | 6,000 | 6,461 | 38 | 52 | 219 | 66 | 8 | 6 |
| Calif. | 2,416 | 3,525 | 89,908 | 90,654 | 180 | 203 | 239 | 220 | - | - |
| Alaska | 17 | 19 | 3,121 | 2,331 | 1 | 1 | 7 | 4 | - | - |
| Hawaii | 66 | 60 | 3,818 | 3,578 | 2 | 3 | 39 | 17 | - | - |
| Guam | 2 | 11 | - | 359 | - | - | N | N | - | - |
| P.R. | 668 | 1,017 | 1,997 | 2,432 | - | - | - | 2 | - | - |
| V.I. | 66 | 2 | 125 | 132 | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | 2 | U | 138 | U | - | U | - | U | - | U |

N: Not notifiable. U: Unavailable. -: No reported cases. C.N.M.I.: Commonwealth of Northern Mariana Islands.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

† Chlamydia refers to genital infections caused by *C. trachomatis*.

‡ Updated monthly from reports to the Division of HIV/AIDS Prevention — Surveillance and Epidemiology, National Center for HIV, STD, and TB Prevention. Last update October 31, 2002.

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Escherichia coli Enterohemorrhagic | | Giardiasis | | | Haemophilus influenzae, Invasive | | | |
|----------------|--|--------------|------------|---------------|--------------|-------------------------------------|--------------|--------------|--------------|
| | Shiga Toxin Positive, Not Serogrouped | | | | | All Ages, All Serotypes | | Age <5 Years | |
| | | | | Serotype B | | | | | |
| | Cum. 2002 | Cum. 2001 | | Cum. 2002 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 |
| UNITED STATES | 39 | 17 | 15,103 | 288,197 | 316,755 | 1,335 | 1,291 | 22 | 22 |
| NEW ENGLAND | 1 | 1 | 1,497 | 6,704 | 6,108 | 115 | 98 | - | 1 |
| Maine | - | - | 192 | 120 | 119 | 1 | 2 | - | - |
| N.H. | - | - | 41 | 116 | 161 | 8 | 6 | - | - |
| Vt. | 1 | 1 | 129 | 84 | 61 | 7 | 3 | - | - |
| Mass. | - | - | 752 | 2,914 | 2,810 | 50 | 41 | - | - |
| R.I. | - | - | 140 | 841 | 753 | 10 | 5 | - | 1 |
| Conn. | - | - | 243 | 2,629 | 2,204 | 39 | 41 | - | - |
| MID. ATLANTIC | - | 3 | 3,264 | 35,186 | 37,697 | 235 | 197 | 4 | 3 |
| Upstate N.Y. | - | - | 1,111 | 7,664 | 7,554 | 106 | 67 | 2 | - |
| N.Y. City | - | - | 1,151 | 10,284 | 11,186 | 55 | 51 | - | - |
| N.J. | - | - | 342 | 6,130 | 7,230 | 49 | 44 | - | - |
| Pa. | - | 3 | 660 | 11,108 | 11,727 | 25 | 35 | 2 | 3 |
| E.N. CENTRAL | 12 | 6 | 2,908 | 58,330 | 66,971 | 188 | 242 | 3 | 2 |
| Ohio | 11 | 6 | 850 | 16,157 | 18,829 | 71 | 65 | - | 1 |
| Ind. | - | - | - | 6,538 | 6,164 | 38 | 46 | 1 | - |
| Ill. | - | - | 672 | 17,254 | 21,186 | 57 | 87 | - | - |
| Mich. | 1 | - | 839 | 13,087 | 15,391 | 14 | 13 | 2 | - |
| Wis. | - | - | 547 | 5,294 | 5,401 | 8 | 31 | - | 1 |
| W.N. CENTRAL | 1 | 3 | 1,809 | 14,652 | 14,871 | 60 | 66 | 1 | 1 |
| Minn. | - | - | 714 | 2,564 | 2,322 | 42 | 36 | 1 | - |
| Iowa | - | - | 279 | 1,117 | 1,184 | 1 | - | - | - |
| Mo. | N | N | 431 | 7,696 | 7,696 | 11 | 18 | - | - |
| N. Dak. | 1 | 3 | 27 | 47 | 42 | - | 7 | - | - |
| S. Dak. | - | - | 66 | 232 | 247 | - | - | - | - |
| Nebr. | - | - | 133 | 713 | 1,048 | 1 | 3 | - | - |
| Kans. | - | - | 159 | 2,283 | 2,332 | 5 | 2 | - | 1 |
| S. ATLANTIC | 1 | - | 2,560 | 74,107 | 81,547 | 329 | 319 | 4 | 1 |
| Del. | - | - | 49 | 1,404 | 1,522 | - | - | - | - |
| Md. | - | - | 104 | 7,760 | 8,127 | 78 | 79 | 2 | - |
| D.C. | - | - | 42 | 2,442 | 2,544 | - | - | - | - |
| Va. | - | - | 276 | 8,297 | 9,506 | 29 | 27 | - | - |
| W. Va. | 1 | - | 53 | 812 | 623 | 15 | 14 | - | - |
| N.C. | - | - | - | 14,208 | 15,008 | 30 | 44 | - | 1 |
| S.C. | - | - | 118 | 6,387 | 9,584 | 12 | 6 | - | - |
| Ga. | - | - | 775 | 14,575 | 15,773 | 84 | 86 | - | - |
| Fla. | - | - | 1,143 | 18,222 | 18,860 | 81 | 63 | 2 | - |
| E.S. CENTRAL | 8 | 3 | 337 | 23,838 | 28,452 | 60 | 68 | 1 | - |
| Ky. | 8 | 3 | - | 3,336 | 3,202 | 5 | 2 | - | - |
| Tenn. | - | - | 161 | 8,291 | 8,602 | 30 | 38 | - | - |
| Ala. | - | - | 176 | 7,118 | 9,692 | 16 | 26 | 1 | - |
| Miss. | - | - | - | 5,093 | 6,956 | 9 | 2 | - | - |
| W.S. CENTRAL | 4 | - | 219 | 42,278 | 46,620 | 58 | 51 | 2 | 2 |
| Ark. | - | - | 150 | 4,027 | 4,171 | 2 | 1 | - | - |
| La. | - | - | 4 | 10,446 | 11,103 | 9 | 9 | - | - |
| Okla. | - | - | 65 | 4,088 | 4,294 | 42 | 39 | - | - |
| Tex. | 4 | - | - | 23,717 | 27,052 | 5 | 2 | 2 | 2 |
| MOUNTAIN | 12 | 1 | 1,483 | 8,869 | 9,160 | 172 | 131 | 4 | 8 |
| Mont. | - | - | 78 | 99 | 94 | - | - | - | - |
| Idaho | - | - | 119 | 83 | 69 | 2 | 2 | - | - |
| Wyo. | - | - | 29 | 55 | 76 | 1 | 1 | - | - |
| Colo. | 12 | 1 | 494 | 3,009 | 2,829 | 31 | 37 | - | - |
| N. Mex. | - | - | 132 | 1,204 | 893 | 25 | 21 | - | - |
| Ariz. | - | - | 190 | 3,220 | 3,460 | 84 | 52 | 2 | 1 |
| Utah | - | - | 298 | 239 | 169 | 17 | 7 | 1 | 4 |
| Nev. | - | - | 143 | 960 | 1,570 | 12 | 11 | 1 | 1 |
| PACIFIC | - | - | 1,026 | 24,233 | 25,329 | 118 | 119 | 3 | 4 |
| Wash. | - | - | 376 | 2,558 | 2,672 | 3 | 5 | 2 | - |
| Oreg. | - | - | 401 | 764 | 995 | 59 | 34 | - | - |
| Calif. | - | - | 73 | 19,765 | 20,732 | 22 | 52 | 1 | 4 |
| Alaska | - | - | 96 | 516 | 381 | 1 | 6 | - | - |
| Hawaii | - | - | 80 | 630 | 549 | 33 | 22 | - | - |
| Guam | - | - | - | - | 45 | - | - | - | - |
| P.R. | - | - | 38 | 292 | 530 | 1 | 1 | - | - |
| V.I. | - | - | - | 31 | 25 | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | 1 | 13 | U | - | U | - | U |

N: Not notifiable.

U: Unavailable.

- : No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Haemophilus influenzae, Invasive | | | | Hepatitis (Viral, Acute), By Type | | | | | |
|----------------|----------------------------------|-----------|------------------|-----------|-----------------------------------|-----------|-----------|-----------|-----------------|-----------|
| | Age <5 Years | | | | | | | | | |
| | Non-Serotype B | | Unknown Serotype | | A | | B | | C: Non-A, Non-B | |
| | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| UNITED STATES | 222 | 216 | 15 | 26 | 7,446 | 9,095 | 6,011 | 6,411 | 3,098 | 3,506 |
| NEW ENGLAND | 13 | 15 | - | - | 266 | 643 | 212 | 125 | 22 | 33 |
| Maine | - | - | - | - | 8 | 11 | 11 | 5 | - | - |
| N.H. | - | 1 | - | - | 11 | 15 | 20 | 13 | - | - |
| Vt. | - | - | - | - | 1 | 16 | 4 | 5 | 13 | 7 |
| Mass. | 8 | 7 | - | - | 129 | 322 | 115 | 30 | 9 | 26 |
| R.I. | - | - | - | - | 30 | 59 | 26 | 25 | - | - |
| Conn. | 5 | 7 | - | - | 87 | 220 | 36 | 47 | - | - |
| MID. ATLANTIC | 28 | 32 | - | 3 | 941 | 1,149 | 1,366 | 1,219 | 1,592 | 1,198 |
| Upstate N.Y. | 12 | 9 | - | 1 | 168 | 239 | 122 | 111 | 62 | 26 |
| N.Y. City | 8 | 11 | - | - | 456 | 399 | 695 | 569 | - | - |
| N.J. | 5 | 5 | - | - | 122 | 262 | 345 | 266 | 1,499 | 1,109 |
| Pa. | 3 | 7 | - | 2 | 195 | 249 | 204 | 273 | 31 | 63 |
| E.N. CENTRAL | 31 | 38 | 1 | 2 | 983 | 1,094 | 561 | 852 | 92 | 150 |
| Ohio | 8 | 12 | 1 | - | 297 | 225 | 94 | 88 | 8 | 8 |
| Ind. | 7 | 6 | - | 1 | 45 | 91 | 43 | 46 | - | 1 |
| Ill. | 11 | 14 | - | - | 252 | 402 | 126 | 132 | 13 | 11 |
| Mich. | 3 | - | - | 1 | 216 | 302 | 298 | 546 | 71 | 130 |
| Wis. | 2 | 6 | - | - | 173 | 74 | - | 40 | - | - |
| W.N. CENTRAL | 6 | 5 | 3 | 6 | 281 | 352 | 202 | 190 | 721 | 1,027 |
| Minn. | 5 | 3 | 1 | 2 | 39 | 40 | 28 | 21 | - | 9 |
| Iowa | - | - | - | - | 74 | 33 | 17 | 21 | 1 | - |
| Mo. | - | - | 2 | 4 | 78 | 77 | 108 | 108 | 702 | 1,005 |
| N. Dak. | - | 1 | - | - | 3 | 3 | 5 | 1 | - | - |
| S. Dak. | - | - | - | - | 3 | 3 | 2 | 1 | 1 | - |
| Nebr. | 1 | 1 | - | - | 17 | 32 | 22 | 26 | 13 | 6 |
| Kans. | - | - | - | - | 67 | 164 | 20 | 12 | 4 | 7 |
| S. ATLANTIC | 45 | 44 | 2 | 6 | 2,151 | 2,207 | 1,461 | 1,349 | 168 | 95 |
| Del. | - | - | - | - | 12 | 16 | 7 | 25 | 5 | 10 |
| Md. | 4 | 8 | - | 1 | 277 | 235 | 109 | 130 | 6 | 8 |
| D.C. | - | - | - | - | 71 | 51 | 22 | 11 | - | - |
| Va. | 4 | 5 | - | - | 132 | 120 | 178 | 158 | 16 | - |
| W. Va. | 1 | 1 | 1 | 1 | 18 | 18 | 18 | 20 | 3 | 9 |
| N.C. | 3 | 2 | - | 4 | 197 | 206 | 207 | 191 | 25 | 19 |
| S.C. | 2 | 1 | - | - | 56 | 70 | 112 | 29 | 4 | 6 |
| Ga. | 18 | 18 | - | - | 410 | 855 | 338 | 388 | 29 | - |
| Fla. | 13 | 9 | 1 | - | 978 | 636 | 470 | 397 | 80 | 43 |
| E.S. CENTRAL | 13 | 12 | 1 | 3 | 241 | 368 | 339 | 424 | 181 | 182 |
| Ky. | 1 | - | - | 1 | 41 | 122 | 48 | 49 | 3 | 9 |
| Tenn. | 7 | 6 | - | 1 | 109 | 142 | 120 | 214 | 25 | 63 |
| Ala. | 3 | 5 | 1 | 1 | 36 | 71 | 95 | 79 | 10 | 4 |
| Miss. | 2 | 1 | - | - | 55 | 33 | 76 | 82 | 143 | 106 |
| W.S. CENTRAL | 14 | 9 | - | - | 554 | 773 | 605 | 759 | 160 | 649 |
| Ark. | 1 | 1 | - | - | 45 | 64 | 79 | 89 | 7 | 10 |
| La. | 2 | 2 | - | - | 65 | 85 | 94 | 112 | 66 | 143 |
| Okla. | 9 | 6 | - | - | 46 | 107 | 44 | 85 | 5 | 4 |
| Tex. | 2 | - | - | - | 398 | 517 | 388 | 473 | 82 | 492 |
| MOUNTAIN | 49 | 21 | 7 | 1 | 518 | 640 | 557 | 412 | 60 | 50 |
| Mont. | - | - | - | - | 13 | 11 | 9 | 3 | 1 | 1 |
| Idaho | 1 | - | - | - | 28 | 52 | 6 | 11 | 1 | 2 |
| Wyo. | - | - | - | - | 3 | 7 | 17 | 3 | 5 | 7 |
| Colo. | 3 | 2 | - | - | 72 | 80 | 70 | 89 | 18 | 8 |
| N. Mex. | 6 | 9 | 1 | 1 | 28 | 40 | 140 | 119 | 1 | 11 |
| Ariz. | 30 | 8 | 5 | - | 265 | 323 | 204 | 122 | 4 | 9 |
| Utah | 5 | 2 | - | - | 62 | 64 | 56 | 22 | 4 | 3 |
| Nev. | 4 | - | 1 | - | 47 | 63 | 55 | 43 | 26 | 9 |
| PACIFIC | 23 | 40 | 1 | 5 | 1,511 | 1,869 | 708 | 1,081 | 102 | 122 |
| Wash. | 1 | 3 | - | 2 | 141 | 138 | 64 | 129 | 24 | 21 |
| Oreg. | 5 | 7 | - | - | 61 | 92 | 113 | 151 | 16 | 14 |
| Calif. | 13 | 28 | 1 | 1 | 1,297 | 1,609 | 519 | 774 | 62 | 87 |
| Alaska | 1 | 1 | - | - | 10 | 14 | 4 | 9 | - | - |
| Hawaii | 3 | 1 | - | 2 | 2 | 16 | 8 | 18 | - | - |
| Guam | - | - | - | - | - | 1 | - | - | - | - |
| P.R. | - | 1 | - | - | 96 | 204 | 84 | 238 | - | 1 |
| V.I. | - | - | - | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | - | U | - | U | 37 | U | - | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Legionellosis | | Listeriosis | | Lyme Disease | | Malaria | | Measles Total | |
|----------------|---------------|-----------|-------------|-----------|--------------|-----------|-----------|-----------|-----------------|------------------|
| | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| UNITED STATES | 1,030 | 981 | 527 | 538 | 15,161 | 13,725 | 1,136 | 1,314 | 36 [†] | 114 [‡] |
| NEW ENGLAND | 94 | 68 | 55 | 52 | 4,426 | 3,973 | 58 | 87 | - | 5 |
| Maine | 3 | 8 | 5 | 2 | 111 | - | 5 | 4 | - | - |
| N.H. | 6 | 10 | 4 | 4 | 232 | 97 | 7 | 2 | - | - |
| Vt. | 36 | 5 | 3 | 3 | 32 | 17 | 4 | 1 | - | 1 |
| Mass. | 30 | 19 | 29 | 27 | 1,150 | 1,110 | 21 | 47 | - | 3 |
| R.I. | 5 | 10 | 1 | 1 | 327 | 449 | 7 | 9 | - | - |
| Conn. | 14 | 14 | 13 | 15 | 2,574 | 2,300 | 14 | 24 | - | 1 |
| MID. ATLANTIC | 282 | 232 | 148 | 95 | 8,869 | 7,538 | 292 | 396 | 7 | 20 |
| Upstate N.Y. | 94 | 61 | 54 | 25 | 4,680 | 3,165 | 43 | 59 | 1 | 4 |
| N.Y. City | 49 | 43 | 30 | 23 | 150 | 61 | 181 | 234 | 6 | 7 |
| N.J. | 27 | 21 | 31 | 17 | 1,641 | 1,980 | 36 | 60 | - | 1 |
| Pa. | 112 | 107 | 33 | 30 | 2,398 | 2,332 | 32 | 43 | - | 8 |
| E.N. CENTRAL | 239 | 281 | 69 | 82 | 86 | 703 | 124 | 159 | 3 | 10 |
| Ohio | 110 | 122 | 24 | 14 | 67 | 40 | 22 | 23 | 1 | 3 |
| Ind. | 20 | 20 | 9 | 8 | 19 | 22 | 12 | 16 | 2 | 4 |
| Ill. | - | 24 | 11 | 23 | - | 31 | 30 | 65 | - | 3 |
| Mich. | 75 | 73 | 18 | 24 | - | 17 | 46 | 36 | - | - |
| Wis. | 34 | 42 | 7 | 13 | U | 593 | 14 | 19 | - | - |
| W.N. CENTRAL | 54 | 46 | 17 | 17 | 360 | 364 | 56 | 34 | 3 | 5 |
| Minn. | 14 | 9 | 3 | - | 269 | 292 | 17 | 6 | 1 | 3 |
| Iowa | 11 | 8 | 2 | 2 | 36 | 34 | 4 | 7 | - | - |
| Mo. | 15 | 20 | 8 | 10 | 40 | 32 | 15 | 13 | 2 | 2 |
| N. Dak. | - | 1 | 1 | - | 1 | - | 1 | - | - | - |
| S. Dak. | 4 | 3 | 1 | - | 2 | - | 1 | - | - | - |
| Nebr. | 10 | 4 | 1 | 1 | 6 | 4 | 5 | 2 | - | - |
| Kans. | - | 1 | 1 | 4 | 6 | 2 | 13 | 6 | - | - |
| S. ATLANTIC | 189 | 168 | 76 | 72 | 1,192 | 893 | 323 | 266 | 2 | 5 |
| Del. | 9 | 12 | - | 2 | 161 | 152 | 4 | 2 | - | - |
| Md. | 42 | 32 | 18 | 14 | 632 | 543 | 104 | 108 | - | 3 |
| D.C. | 6 | 8 | - | - | 21 | 15 | 19 | 13 | - | - |
| Va. | 24 | 20 | 7 | 12 | 145 | 115 | 31 | 45 | - | 1 |
| W. Va. | N | N | - | 5 | 17 | 11 | 3 | 1 | - | - |
| N.C. | 11 | 10 | 6 | 5 | 124 | 38 | 21 | 17 | - | - |
| S.C. | 8 | 13 | 8 | 5 | 20 | 5 | 7 | 6 | - | - |
| Ga. | 17 | 11 | 12 | 14 | 2 | - | 73 | 42 | - | 1 |
| Fla. | 72 | 62 | 25 | 15 | 70 | 14 | 61 | 32 | 2 | - |
| E.S. CENTRAL | 41 | 56 | 19 | 22 | 47 | 63 | 19 | 35 | 12 | 2 |
| Ky. | 19 | 12 | 4 | 7 | 22 | 23 | 7 | 14 | - | 2 |
| Tenn. | 14 | 27 | 11 | 8 | 22 | 25 | 3 | 11 | - | - |
| Ala. | 8 | 13 | 4 | 7 | 3 | 8 | 4 | 6 | 12 | - |
| Miss. | - | 4 | - | - | - | 7 | 5 | 4 | - | - |
| W.S. CENTRAL | 16 | 24 | 18 | 31 | 17 | 82 | 16 | 83 | 2 | 1 |
| Ark. | - | - | - | 1 | 3 | - | 2 | 3 | - | - |
| La. | 4 | 6 | - | - | 4 | 8 | 4 | 6 | - | - |
| Okla. | 3 | 3 | 9 | 2 | - | - | 9 | 3 | - | - |
| Tex. | 9 | 15 | 9 | 28 | 10 | 74 | 1 | 71 | 2 | 1 |
| MOUNTAIN | 47 | 49 | 27 | 36 | 20 | 11 | 45 | 54 | 2 | 2 |
| Mont. | 3 | - | - | - | - | - | 2 | 3 | - | - |
| Idaho | 1 | 3 | 2 | 1 | 4 | 5 | - | 3 | - | 1 |
| Wyo. | 1 | 2 | - | 2 | 1 | - | - | - | - | - |
| Colo. | 7 | 14 | 6 | 9 | 3 | 1 | 22 | 22 | - | - |
| N. Mex. | 2 | 3 | 3 | 7 | 1 | - | 3 | 3 | - | - |
| Ariz. | 13 | 16 | 12 | 8 | 3 | - | 10 | 10 | - | 1 |
| Utah | 15 | 7 | 3 | 2 | 7 | 1 | 5 | 4 | 1 | - |
| Nev. | 5 | 4 | 1 | 7 | 1 | 3 | 3 | 9 | 1 | - |
| PACIFIC | 68 | 59 | 98 | 131 | 144 | 98 | 203 | 200 | 5 | 64 |
| Wash. | 7 | 9 | 8 | 10 | 10 | 7 | 22 | 10 | - | 15 |
| Oreg. | N | N | 9 | 12 | 15 | 11 | 9 | 16 | - | 3 |
| Calif. | 60 | 44 | 73 | 103 | 116 | 78 | 163 | 162 | 3 | 39 |
| Alaska | - | 1 | - | - | 3 | 2 | 2 | 1 | - | - |
| Hawaii | 1 | 5 | 8 | 6 | N | N | 7 | 11 | 2 | 7 |
| Guam | - | - | - | - | - | - | - | 1 | - | - |
| P.R. | - | 2 | 1 | - | N | N | - | 5 | - | 1 |
| V.I. | - | - | - | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | - | U | - | U | - | U | - | U |

N: Not notifiable.

U: Unavailable.

-: No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

† Of 36 cases reported, 23 were indigenous and 13 were imported from another country.

‡ Of 114 cases reported, 60 were indigenous and 54 were imported from another country.

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Meningococcal Disease | | Mumps | | Pertussis | | Rabies, Animal | |
|----------------|-----------------------|-----------|-----------|-----------|-----------|-----------|----------------|-----------|
| | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| UNITED STATES | 1,467 | 2,031 | 236 | 217 | 6,632 | 4,939 | 5,507 | 6,418 |
| NEW ENGLAND | 80 | 95 | 8 | 2 | 552 | 496 | 841 | 663 |
| Maine | 7 | 4 | 1 | - | 17 | 22 | 54 | 63 |
| N.H. | 11 | 12 | 4 | - | 17 | 27 | 45 | 19 |
| Vt. | 4 | 5 | - | - | 124 | 35 | 87 | 58 |
| Mass. | 39 | 52 | 2 | 2 | 355 | 389 | 276 | 246 |
| R.I. | 5 | 4 | - | - | 13 | 5 | 70 | 65 |
| Conn. | 14 | 18 | 1 | - | 26 | 18 | 309 | 212 |
| MID. ATLANTIC | 136 | 231 | 24 | 25 | 411 | 320 | 1,037 | 1,187 |
| Upstate N.Y. | 40 | 64 | 6 | 3 | 298 | 130 | 637 | 724 |
| N.Y. City | 22 | 41 | 2 | 12 | 13 | 53 | 10 | 36 |
| N.J. | 26 | 40 | - | 3 | 4 | 18 | 171 | 175 |
| Pa. | 48 | 86 | 16 | 7 | 96 | 119 | 219 | 252 |
| E.N. CENTRAL | 194 | 315 | 38 | 27 | 801 | 767 | 147 | 148 |
| Ohio | 72 | 80 | 13 | 1 | 380 | 280 | 39 | 42 |
| Ind. | 31 | 35 | 2 | 3 | 123 | 79 | 31 | 15 |
| Ill. | 36 | 79 | 14 | 16 | 144 | 90 | 31 | 24 |
| Mich. | 40 | 73 | 8 | 5 | 49 | 133 | 46 | 47 |
| Wis. | 15 | 48 | 1 | 2 | 105 | 185 | - | 20 |
| W.N. CENTRAL | 137 | 141 | 16 | 8 | 673 | 344 | 407 | 342 |
| Minn. | 32 | 20 | 4 | 3 | 340 | 146 | 36 | 43 |
| Iowa | 21 | 29 | 1 | - | 130 | 65 | 72 | 77 |
| Mo. | 45 | 50 | 5 | 1 | 129 | 93 | 49 | 40 |
| N. Dak. | 3 | 6 | 1 | - | 1 | 4 | 29 | 35 |
| S. Dak. | 2 | 5 | - | - | 6 | 4 | 65 | 55 |
| Nebr. | 26 | 17 | - | 1 | 8 | 5 | - | 4 |
| Kans. | 8 | 14 | 5 | 3 | 59 | 27 | 156 | 88 |
| S. ATLANTIC | 263 | 316 | 25 | 38 | 376 | 230 | 2,266 | 2,247 |
| Del. | 7 | 5 | - | - | 3 | - | 24 | 30 |
| Md. | 8 | 38 | 5 | 6 | 57 | 42 | 321 | 466 |
| D.C. | - | - | - | - | 2 | 1 | - | - |
| Va. | 40 | 37 | 4 | 8 | 133 | 40 | 460 | 441 |
| W. Va. | 4 | 12 | - | - | 31 | 3 | 161 | 131 |
| N.C. | 30 | 62 | 2 | 5 | 40 | 69 | 654 | 522 |
| S.C. | 28 | 31 | 3 | 5 | 41 | 31 | 133 | 102 |
| Ga. | 34 | 48 | 4 | 9 | 21 | 21 | 347 | 372 |
| Fla. | 112 | 83 | 7 | 5 | 48 | 23 | 166 | 183 |
| E.S. CENTRAL | 86 | 125 | 13 | 9 | 236 | 177 | 156 | 197 |
| Ky. | 14 | 22 | 3 | 3 | 91 | 80 | 26 | 26 |
| Tenn. | 36 | 56 | 2 | 1 | 104 | 58 | 98 | 106 |
| Ala. | 22 | 31 | 3 | - | 32 | 35 | 28 | 61 |
| Miss. | 14 | 16 | 5 | 5 | 9 | 4 | 4 | 4 |
| W.S. CENTRAL | 180 | 298 | 17 | 13 | 1,485 | 594 | 112 | 1,030 |
| Ark. | 23 | 21 | - | - | 468 | 151 | 3 | - |
| La. | 33 | 74 | 1 | 2 | 7 | 8 | - | 8 |
| Okla. | 20 | 28 | - | - | 66 | 28 | 108 | 57 |
| Tex. | 104 | 175 | 16 | 11 | 944 | 407 | 1 | 965 |
| MOUNTAIN | 86 | 87 | 17 | 14 | 938 | 1,225 | 279 | 253 |
| Mont. | 2 | 4 | - | 1 | 5 | 36 | 18 | 38 |
| Idaho | 4 | 7 | 1 | 1 | 65 | 170 | 38 | 28 |
| Wyo. | - | 5 | - | 1 | 11 | 1 | 18 | 28 |
| Colo. | 21 | 34 | 2 | 3 | 374 | 294 | 59 | - |
| N. Mex. | 4 | 10 | 1 | 2 | 170 | 129 | 7 | 15 |
| Ariz. | 30 | 13 | 1 | 1 | 169 | 496 | 115 | 128 |
| Utah | 5 | 8 | 7 | 1 | 97 | 75 | 13 | 15 |
| Nev. | 20 | 6 | 5 | 4 | 47 | 24 | 11 | 1 |
| PACIFIC | 305 | 423 | 78 | 81 | 1,160 | 786 | 262 | 351 |
| Wash. | 60 | 59 | - | 2 | 399 | 140 | - | - |
| Oreg. | 42 | 56 | N | N | 173 | 51 | 13 | 4 |
| Calif. | 191 | 293 | 64 | 39 | 567 | 549 | 225 | 309 |
| Alaska | 4 | 2 | - | 1 | 4 | 11 | 24 | 38 |
| Hawaii | 8 | 13 | 14 | 39 | 17 | 35 | - | - |
| Guam | - | - | - | - | - | - | - | - |
| P.R. | 5 | 5 | - | 1 | 3 | - | 49 | 85 |
| V.I. | - | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | - | U | 1 | U | - | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Rocky Mountain Spotted Fever | | Rubella | | | | Salmonellosis | |
|----------------|---------------------------------|--------------|--------------|--------------|-----------------------|--------------|---------------|--------------|
| | | | Rubella | | Congenital Rubella | | | |
| | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| UNITED STATES | 934 | 538 | 13 | 21 | 2 | - | 36,774 | 35,540 |
| NEW ENGLAND | 8 | 3 | - | - | - | - | 1,994 | 2,158 |
| Maine | - | - | - | - | - | - | 134 | 161 |
| N.H. | - | 1 | - | - | - | - | 126 | 155 |
| Vt. | - | - | - | - | - | - | 70 | 74 |
| Mass. | 4 | 2 | - | - | - | - | 1,103 | 1,245 |
| R.I. | 4 | - | - | - | - | - | 153 | 120 |
| Conn. | - | - | - | - | - | - | 408 | 403 |
| MID. ATLANTIC | 42 | 31 | 1 | 8 | - | - | 4,596 | 4,655 |
| Upstate N.Y. | 7 | 2 | 1 | 1 | - | - | 1,402 | 1,077 |
| N.Y. City | 9 | 2 | - | 6 | - | - | 1,278 | 1,174 |
| N.J. | 10 | 9 | - | 1 | - | - | 671 | 1,084 |
| Pa. | 16 | 18 | - | - | - | - | 1,245 | 1,320 |
| E.N. CENTRAL | 18 | 16 | 1 | 2 | - | - | 4,749 | 4,531 |
| Ohio | 12 | 2 | - | - | - | - | 1,285 | 1,227 |
| Ind. | 3 | 1 | - | - | - | - | 441 | 478 |
| Ill. | - | 12 | - | 2 | - | - | 1,445 | 1,266 |
| Mich. | 3 | 1 | 1 | - | - | - | 804 | 794 |
| Wis. | - | - | - | - | - | - | 774 | 766 |
| W.N. CENTRAL | 97 | 67 | - | 3 | - | - | 2,375 | 2,068 |
| Minn. | - | - | - | - | - | - | 523 | 558 |
| Iowa | 3 | 2 | - | 1 | - | - | 461 | 320 |
| Mo. | 89 | 61 | - | 1 | - | - | 791 | 567 |
| N. Dak. | - | 1 | - | - | - | - | 42 | 58 |
| S. Dak. | 1 | 2 | - | - | - | - | 102 | 143 |
| Nebr. | 4 | 1 | - | - | - | - | 150 | 143 |
| Kans. | - | - | - | 1 | - | - | 306 | 279 |
| S. ATLANTIC | 497 | 264 | 5 | 5 | - | - | 10,167 | 8,346 |
| Del. | 4 | 10 | - | - | - | - | 87 | 90 |
| Md. | 56 | 38 | - | 1 | - | - | 854 | 711 |
| D.C. | 2 | - | - | - | - | - | 71 | 75 |
| Va. | 39 | 25 | - | - | - | - | 1,112 | 1,205 |
| W. Va. | 2 | - | - | - | - | - | 137 | 126 |
| N.C. | 274 | 149 | - | - | - | - | 1,367 | 1,209 |
| S.C. | 68 | 29 | - | 2 | - | - | 720 | 792 |
| Ga. | 27 | 9 | - | - | - | - | 1,855 | 1,539 |
| Fla. | 15 | 4 | 5 | 2 | - | - | 3,964 | 2,599 |
| E.S. CENTRAL | 102 | 106 | - | - | 1 | - | 2,873 | 2,467 |
| Ky. | 5 | 2 | - | - | - | - | 353 | 337 |
| Tenn. | 76 | 74 | - | - | 1 | - | 738 | 583 |
| Ala. | 18 | 15 | - | - | - | - | 806 | 692 |
| Miss. | 3 | 15 | - | - | - | - | 976 | 855 |
| W.S. CENTRAL | 159 | 39 | 2 | 1 | - | - | 3,262 | 4,633 |
| Ark. | 97 | 8 | - | - | - | - | 986 | 842 |
| La. | - | 2 | - | - | - | - | 725 | 792 |
| Okla. | 61 | 29 | - | - | - | - | 457 | 438 |
| Tex. | 1 | - | 2 | 1 | - | - | 1,094 | 2,561 |
| MOUNTAIN | 14 | 11 | 1 | - | - | - | 1,978 | 1,955 |
| Mont. | 1 | 1 | - | - | - | - | 81 | 71 |
| Idaho | - | 1 | - | - | - | - | 135 | 127 |
| Wyo. | 5 | 2 | - | - | - | - | 92 | 58 |
| Colo. | 2 | 2 | - | - | - | - | 496 | 542 |
| N. Mex. | 1 | 1 | - | - | - | - | 281 | 263 |
| Ariz. | - | - | - | - | - | - | 528 | 534 |
| Utah | - | 3 | 1 | - | - | - | 187 | 199 |
| Nev. | 5 | 1 | - | - | - | - | 178 | 161 |
| PACIFIC | 7 | 1 | 3 | 2 | 1 | - | 4,780 | 4,727 |
| Wash. | - | - | - | - | - | - | 471 | 464 |
| Oreg. | 2 | 1 | - | - | - | - | 325 | 250 |
| Calif. | 5 | - | 3 | 1 | - | - | 3,660 | 3,650 |
| Alaska | - | - | - | - | - | - | 72 | 41 |
| Hawaii | - | - | - | 1 | 1 | - | 252 | 322 |
| Guam | - | - | - | - | - | - | - | 24 |
| P.R. | - | - | - | 3 | - | - | 201 | 834 |
| V.I. | - | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | - | U | - | U | - | U | 25 | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Shigellosis | | Streptococcal Disease, Invasive, Group A | | Streptococcus pneumoniae, Drug Resistant, Invasive | | Streptococcus pneumoniae, Invasive (<5 Years) | |
|----------------|-------------|-----------|--|-----------|--|-----------|---|-----------|
| | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| UNITED STATES | 16,530 | 17,183 | 3,624 | 3,249 | 2,058 | 2,288 | 235 | 374 |
| NEW ENGLAND | 296 | 280 | 168 | 204 | 18 | 114 | 3 | 40 |
| Maine | 10 | 6 | 20 | 12 | - | - | - | - |
| N.H. | 11 | 6 | 35 | N | - | - | N | N |
| Vt. | 1 | 7 | 9 | 15 | 5 | 7 | 2 | 1 |
| Mass. | 176 | 197 | 89 | 60 | N | N | N | N |
| R.I. | 17 | 17 | 15 | 12 | 13 | 4 | 1 | 3 |
| Conn. | 81 | 47 | - | 105 | - | 103 | - | 36 |
| MID. ATLANTIC | 1,251 | 1,372 | 590 | 601 | 102 | 147 | 61 | 97 |
| Upstate N.Y. | 291 | 440 | 265 | 239 | 84 | 140 | 59 | 97 |
| N.Y. City | 386 | 379 | 135 | 157 | U | U | U | U |
| N.J. | 349 | 255 | 128 | 128 | N | N | N | N |
| Pa. | 225 | 298 | 62 | 77 | 18 | 7 | 2 | - |
| E.N. CENTRAL | 1,592 | 3,960 | 658 | 721 | 209 | 166 | 104 | 116 |
| Ohio | 575 | 2,633 | 198 | 184 | 60 | 3 | 21 | - |
| Ind. | 91 | 205 | 45 | 56 | 144 | 163 | 57 | 54 |
| Ill. | 611 | 554 | 145 | 232 | 2 | - | - | 62 |
| Mich. | 167 | 283 | 269 | 198 | 3 | - | N | N |
| Wis. | 148 | 285 | 1 | 51 | N | N | 26 | - |
| W.N. CENTRAL | 922 | 1,760 | 224 | 343 | 417 | 139 | 49 | 54 |
| Minn. | 205 | 394 | 113 | 156 | 292 | 63 | 49 | 45 |
| Iowa | 115 | 347 | - | - | N | N | N | N |
| Mo. | 171 | 289 | 42 | 70 | 5 | 9 | - | - |
| N. Dak. | 16 | 21 | 3 | 17 | 1 | 6 | - | 9 |
| S. Dak. | 150 | 554 | 13 | 11 | 1 | 4 | - | - |
| Nebr. | 179 | 87 | 18 | 39 | 29 | 19 | N | N |
| Kans. | 86 | 68 | 35 | 50 | 89 | 38 | N | N |
| S. ATLANTIC | 6,109 | 2,489 | 723 | 531 | 1,078 | 1,203 | 8 | 6 |
| Del. | 296 | 14 | 2 | 4 | 3 | 6 | N | N |
| Md. | 1,068 | 137 | 126 | N | N | N | N | N |
| D.C. | 56 | 54 | 7 | 21 | 49 | 5 | 1 | 4 |
| Va. | 892 | 365 | 68 | 70 | N | N | N | N |
| W. Va. | 12 | 8 | 19 | 19 | 42 | 37 | 7 | 2 |
| N.C. | 399 | 313 | 112 | 134 | N | N | U | U |
| S.C. | 106 | 237 | 34 | 11 | 169 | 247 | N | N |
| Ga. | 1,393 | 479 | 154 | 167 | 270 | 378 | N | N |
| Fla. | 1,887 | 882 | 201 | 105 | 545 | 530 | N | N |
| E.S. CENTRAL | 1,313 | 1,548 | 104 | 108 | 120 | 218 | - | - |
| Ky. | 165 | 733 | 18 | 36 | 17 | 24 | N | N |
| Tenn. | 99 | 92 | 86 | 72 | 103 | 193 | N | N |
| Ala. | 737 | 196 | - | - | - | 1 | N | N |
| Miss. | 312 | 527 | - | - | - | - | - | - |
| W.S. CENTRAL | 1,608 | 2,651 | 105 | 296 | 74 | 259 | 6 | 61 |
| Ark. | 184 | 540 | 7 | - | 7 | 16 | - | - |
| La. | 390 | 223 | - | 1 | 67 | 243 | 3 | 61 |
| Okla. | 525 | 81 | 41 | 39 | N | N | 3 | - |
| Tex. | 509 | 1,807 | 57 | 256 | N | N | - | - |
| MOUNTAIN | 810 | 872 | 529 | 369 | 40 | 38 | 4 | - |
| Mont. | 3 | 8 | - | - | - | - | - | - |
| Idaho | 15 | 39 | 9 | 7 | N | N | N | N |
| Wyo. | 9 | 7 | 7 | 11 | 9 | 7 | - | - |
| Colo. | 163 | 226 | 133 | 143 | - | - | - | - |
| N. Mex. | 194 | 112 | 96 | 77 | 30 | 29 | - | - |
| Ariz. | 347 | 356 | 254 | 128 | - | - | N | N |
| Utah | 34 | 55 | 30 | 3 | - | - | 4 | - |
| Nev. | 45 | 69 | - | - | 1 | 2 | - | - |
| PACIFIC | 2,629 | 2,251 | 523 | 76 | - | 4 | - | - |
| Wash. | 158 | 187 | 65 | - | - | - | N | N |
| Oreg. | 103 | 101 | N | N | N | N | N | N |
| Calif. | 2,301 | 1,901 | 366 | - | N | N | N | N |
| Alaska | 6 | 7 | - | - | - | - | N | N |
| Hawaii | 61 | 55 | 92 | 76 | - | 4 | - | - |
| Guam | - | 47 | - | 1 | - | - | - | - |
| P.R. | 8 | 16 | N | N | - | - | N | N |
| V.I. | - | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | - | - | U | U |
| C.N.M.I. | 17 | U | - | U | - | - | - | U |

N: Not notifiable. U: Unavailable. -: No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

TABLE II. (Continued) Provisional cases of selected notifiable diseases, United States, weeks ending November 16, 2002, and November 17, 2001 (46th Week)*

| Reporting Area | Syphilis | | | | Tuberculosis | | Typhoid Fever | |
|----------------|---------------------|-----------|------------|-----------|--------------|-----------|---------------|-----------|
| | Primary & Secondary | | Congenital | | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 |
| | Cum. 2002 | Cum. 2001 | Cum. 2002 | Cum. 2001 | | | | |
| UNITED STATES | 5,583 | 5,339 | 304 | 437 | 10,359 | 12,205 | 238 | 321 |
| NEW ENGLAND | 121 | 56 | - | 8 | 341 | 409 | 14 | 16 |
| Maine | 2 | 1 | - | 3 | 10 | 16 | - | 1 |
| N.H. | 7 | 1 | - | - | 13 | 16 | - | 2 |
| Vt. | 1 | 3 | - | - | - | 4 | - | - |
| Mass. | 81 | 32 | - | 3 | 197 | 211 | 8 | 10 |
| R.I. | 6 | 9 | - | - | 34 | 57 | - | - |
| Conn. | 24 | 10 | - | 2 | 87 | 105 | 6 | 3 |
| MID. ATLANTIC | 628 | 465 | 58 | 70 | 1,859 | 2,011 | 54 | 106 |
| Upstate N.Y. | 29 | 18 | 9 | 5 | 261 | 326 | 9 | 15 |
| N.Y. City | 387 | 253 | 22 | 32 | 933 | 1,000 | 26 | 44 |
| N.J. | 138 | 112 | 26 | 33 | 439 | 438 | 15 | 38 |
| Pa. | 74 | 82 | 1 | - | 226 | 247 | 4 | 9 |
| E.N. CENTRAL | 954 | 949 | 52 | 62 | 1,000 | 1,243 | 18 | 32 |
| Ohio | 147 | 69 | 4 | 2 | 126 | 245 | 6 | 4 |
| Ind. | 61 | 141 | 1 | 10 | 108 | 90 | 2 | 2 |
| Ill. | 287 | 342 | 30 | 40 | 508 | 575 | 1 | 17 |
| Mich. | 435 | 374 | 17 | 6 | 217 | 262 | 4 | 5 |
| Wis. | 24 | 23 | - | 4 | 41 | 71 | 5 | 4 |
| W.N. CENTRAL | 94 | 89 | - | 9 | 471 | 478 | 8 | 15 |
| Minn. | 48 | 31 | - | 2 | 203 | 201 | 3 | 6 |
| Iowa | 2 | 4 | - | - | 24 | 34 | - | - |
| Mo. | 24 | 23 | - | 5 | 117 | 123 | 1 | 9 |
| N. Dak. | - | - | - | - | 4 | 3 | - | - |
| S. Dak. | - | - | - | - | 10 | 12 | - | - |
| Nebr. | 3 | 8 | - | - | 23 | 32 | 4 | - |
| Kans. | 17 | 23 | - | 2 | 90 | 73 | - | - |
| S. ATLANTIC | 1,489 | 1,798 | 67 | 102 | 2,113 | 2,256 | 44 | 41 |
| Del. | 11 | 14 | - | - | 15 | 15 | - | 1 |
| Md. | 172 | 240 | 14 | 4 | 254 | 204 | 7 | 10 |
| D.C. | 58 | 35 | 1 | 2 | - | 51 | - | - |
| Va. | 59 | 93 | 1 | 5 | 166 | 232 | 7 | 11 |
| W. Va. | 2 | 4 | - | - | 28 | 26 | - | - |
| N.C. | 261 | 409 | 18 | 12 | 309 | 307 | 2 | 2 |
| S.C. | 114 | 216 | 8 | 21 | 146 | 150 | - | - |
| Ga. | 315 | 348 | 10 | 22 | 353 | 425 | 9 | 9 |
| Fla. | 497 | 439 | 15 | 36 | 842 | 846 | 19 | 8 |
| E.S. CENTRAL | 417 | 591 | 19 | 30 | 639 | 743 | 4 | 1 |
| Ky. | 83 | 44 | 3 | 1 | 118 | 115 | 4 | - |
| Tenn. | 153 | 294 | 9 | 17 | 253 | 269 | - | 1 |
| Ala. | 140 | 119 | 4 | 5 | 179 | 239 | - | - |
| Miss. | 41 | 134 | 3 | 7 | 89 | 120 | - | - |
| W.S. CENTRAL | 754 | 660 | 64 | 73 | 1,455 | 1,856 | 5 | 18 |
| Ark. | 32 | 33 | 2 | 7 | 112 | 136 | - | - |
| La. | 135 | 159 | - | - | - | 114 | - | - |
| Okla. | 61 | 56 | 3 | 6 | 122 | 136 | 2 | - |
| Tex. | 526 | 412 | 59 | 60 | 1,221 | 1,470 | 3 | 18 |
| MOUNTAIN | 257 | 197 | 15 | 29 | 299 | 493 | 10 | 8 |
| Mont. | - | - | - | - | 6 | 14 | - | 1 |
| Idaho | 5 | 1 | - | - | 9 | 7 | - | - |
| Wyo. | - | 1 | - | - | 3 | 3 | - | - |
| Colo. | 44 | 20 | 1 | 1 | 48 | 116 | 5 | 1 |
| N. Mex. | 30 | 16 | - | 2 | 22 | 49 | 1 | - |
| Ariz. | 162 | 142 | 14 | 26 | 171 | 198 | - | 1 |
| Utah | 8 | 10 | - | - | 26 | 33 | 2 | 1 |
| Nev. | 8 | 7 | - | - | 14 | 73 | 2 | 4 |
| PACIFIC | 869 | 534 | 29 | 54 | 2,182 | 2,716 | 81 | 84 |
| Wash. | 54 | 42 | 1 | - | 198 | 210 | 6 | 5 |
| Oreg. | 20 | 13 | 1 | - | 97 | 98 | 2 | 7 |
| Calif. | 787 | 467 | 26 | 54 | 1,717 | 2,234 | 68 | 68 |
| Alaska | - | - | - | - | 43 | 46 | - | 1 |
| Hawaii | 8 | 12 | 1 | - | 127 | 128 | 5 | 3 |
| Guam | - | 9 | - | 1 | - | 54 | - | 3 |
| P.R. | 227 | 246 | 15 | 13 | 75 | 95 | - | - |
| V.I. | 1 | - | - | - | - | - | - | - |
| Amer. Samoa | U | U | U | U | U | U | U | U |
| C.N.M.I. | 15 | U | - | U | 32 | U | - | U |

N: Not notifiable.

U: Unavailable.

- : No reported cases.

* Incidence data for reporting year 2001 and 2002 are provisional and cumulative (year-to-date).

TABLE III. Deaths in 122 U.S. cities,* week ending November 16, 2002 (46th Week)

| All Causes, By Age (Years) | | | | | | | | All Causes, By Age (Years) | | | | | | | |
|------------------------------|----------|-------|-------|-------|------|----|------------------------|----------------------------|----------|-----|-------|-------|------|----|------------------------|
| Reporting Area | All Ages | >65 | 45-64 | 25-44 | 1-24 | <1 | P&I [†] Total | Reporting Area | All Ages | >65 | 45-64 | 25-44 | 1-24 | <1 | P&I [†] Total |
| NEW ENGLAND | 652 | 464 | 115 | 45 | 12 | 16 | 67 | S. ATLANTIC | 1,097 | 693 | 252 | 100 | 25 | 27 | 75 |
| Boston, Mass. | 151 | 98 | 29 | 11 | 6 | 7 | 17 | Atlanta, Ga. | U | U | U | U | U | U | U |
| Bridgeport, Conn. | 40 | 28 | 5 | 3 | - | 4 | 5 | Baltimore, Md. | 190 | 114 | 50 | 22 | 3 | 1 | 17 |
| Cambridge, Mass. | 26 | 22 | 3 | 1 | - | - | 2 | Charlotte, N.C. | 98 | 55 | 29 | 8 | 2 | 4 | 13 |
| Fall River, Mass. | 38 | 30 | 6 | 2 | - | - | 8 | Jacksonville, Fla. | 137 | 95 | 29 | 5 | 5 | 3 | 7 |
| Hartford, Conn. | 98 | 66 | 22 | 7 | 3 | - | 10 | Miami, Fla. | 105 | 64 | 24 | 12 | 3 | 2 | 9 |
| Lowell, Mass. | 23 | 19 | 1 | 3 | - | - | - | Norfolk, Va. | 41 | 26 | 8 | 2 | 2 | 3 | 2 |
| Lynn, Mass. | 5 | 4 | 1 | - | - | - | 2 | Richmond, Va. | 63 | 41 | 12 | 7 | 2 | 1 | 3 |
| New Bedford, Mass. | 21 | 16 | 2 | 2 | 1 | - | 2 | Savannah, Ga. | 59 | 44 | 7 | 4 | 1 | 3 | 8 |
| New Haven, Conn. | 38 | 21 | 9 | 5 | - | 3 | 6 | St. Petersburg, Fla. | 80 | 52 | 17 | 9 | - | 2 | 3 |
| Providence, R.I. | 68 | 44 | 19 | 3 | 1 | 1 | - | Tampa, Fla. | 161 | 109 | 34 | 13 | 3 | 2 | 8 |
| Somerville, Mass. | 4 | 4 | - | - | - | - | - | Washington, D.C. | 148 | 80 | 41 | 17 | 4 | 6 | 2 |
| Springfield, Mass. | 39 | 31 | 6 | 2 | - | - | 6 | Wilmington, Del. | 15 | 13 | 1 | 1 | - | - | 3 |
| Waterbury, Conn. | 39 | 32 | 3 | 4 | - | - | 2 | | | | | | | | |
| Worcester, Mass. | 62 | 49 | 9 | 2 | 1 | 1 | 7 | E.S. CENTRAL | 792 | 502 | 186 | 59 | 26 | 19 | 65 |
| MID. ATLANTIC | 2,249 | 1,604 | 426 | 147 | 41 | 31 | 103 | Birmingham, Ala. | 119 | 76 | 30 | 8 | 2 | 3 | 6 |
| Albany, N.Y. | 56 | 43 | 6 | 3 | 3 | 1 | 1 | Chattanooga, Tenn. | 57 | 39 | 16 | 1 | - | 1 | 7 |
| Allentown, Pa. | 25 | 23 | 2 | - | - | - | 1 | Knoxville, Tenn. | 80 | 56 | 18 | 3 | 3 | - | 2 |
| Buffalo, N.Y. | 105 | 71 | 24 | 3 | 6 | 1 | 11 | Lexington, Ky. | 59 | 39 | 10 | 6 | 3 | 1 | 2 |
| Camden, N.J. | 33 | 18 | 7 | 7 | - | 1 | 1 | Memphis, Tenn. | 206 | 125 | 48 | 18 | 7 | 8 | 27 |
| Elizabeth, N.J. | 15 | 12 | 2 | 1 | - | - | - | Mobile, Ala. | 54 | 35 | 15 | 2 | 2 | - | 1 |
| Erie, Pa. | 38 | 29 | 5 | 1 | 1 | 2 | 2 | Montgomery, Ala. | 60 | 39 | 10 | 5 | 5 | 1 | 7 |
| Jersey City, N.J. | 48 | 27 | 12 | 7 | - | 2 | - | Nashville, Tenn. | 157 | 93 | 39 | 16 | 4 | 5 | 13 |
| New York City, N.Y. | 1,227 | 880 | 236 | 74 | 23 | 14 | 43 | W.S. CENTRAL | 1,085 | 710 | 243 | 83 | 23 | 18 | 63 |
| Newark, N.J. | 46 | 20 | 16 | 7 | 2 | 1 | 5 | Austin, Tex. | 73 | 51 | 16 | 6 | - | - | 4 |
| Paterson, N.J. | 25 | 13 | 5 | 5 | - | 2 | - | Baton Rouge, La. | 49 | 42 | 6 | - | - | 1 | 6 |
| Philadelphia, Pa. | 230 | 148 | 61 | 16 | 2 | 3 | 3 | Corpus Christi, Tex. | 50 | 36 | 11 | 1 | 1 | 1 | 6 |
| Pittsburgh, Pa. [‡] | 39 | 30 | 7 | 2 | - | - | 3 | Dallas, Tex. | 182 | 103 | 47 | 21 | 6 | 5 | 8 |
| Reading, Pa. | 28 | 24 | 2 | 2 | - | - | 2 | El Paso, Tex. | 121 | 89 | 19 | 11 | 1 | 1 | 7 |
| Rochester, N.Y. | 125 | 105 | 12 | 5 | 1 | 2 | 16 | Fl. Worth, Tex. | 116 | 76 | 25 | 7 | 5 | 3 | 7 |
| Schenectady, N.Y. | 23 | 15 | 3 | 3 | 1 | 1 | 2 | Houston, Tex. | U | U | U | U | U | U | U |
| Scranton, Pa. | 23 | 21 | 1 | 1 | - | - | 2 | Little Rock, Ark. | 53 | 32 | 7 | 4 | - | 2 | - |
| Syracuse, N.Y. | 94 | 74 | 15 | 4 | 1 | - | 7 | New Orleans, La. | U | U | U | U | U | U | U |
| Trenton, N.J. | 22 | 14 | 4 | 2 | 1 | 1 | - | San Antonio, Tex. | 275 | 177 | 66 | 25 | 5 | 2 | 16 |
| Utica, N.Y. | 20 | 17 | 1 | 2 | - | - | 2 | Shreveport, La. | 50 | 30 | 15 | 2 | 2 | 1 | 1 |
| Yonkers, N.Y. | 27 | 20 | 5 | 2 | - | - | 2 | Tulsa, Okla. | 116 | 74 | 31 | 6 | 3 | 2 | 8 |
| E.N. CENTRAL | 1,752 | 1,240 | 333 | 89 | 48 | 42 | 119 | MOUNTAIN | 863 | 562 | 195 | 60 | 25 | 21 | 55 |
| Akron, Ohio | 53 | 42 | 10 | 1 | - | - | 5 | Albuquerque, N.M. | 120 | 79 | 29 | 5 | 5 | 2 | 7 |
| Canton, Ohio | 36 | 24 | 9 | - | - | 3 | 2 | Boise, Idaho | 57 | 35 | 12 | 8 | 1 | 1 | 1 |
| Chicago, Ill. | U | U | U | U | U | U | U | Colo. Springs, Colo. | 72 | 50 | 12 | 6 | 2 | 2 | 3 |
| Cincinnati, Ohio | 63 | 42 | 14 | 3 | 2 | 2 | 8 | Denver, Colo. | 102 | 60 | 25 | 6 | 4 | 7 | 5 |
| Cleveland, Ohio | 124 | 86 | 27 | 8 | 2 | 1 | 5 | Las Vegas, Nev. | 225 | 134 | 64 | 18 | 6 | 3 | 10 |
| Columbus, Ohio | 233 | 158 | 50 | 8 | 9 | 8 | 21 | Ogden, Utah | 31 | 27 | 3 | 1 | - | - | 3 |
| Dayton, Ohio | 121 | 89 | 23 | 6 | 3 | - | 12 | Phoenix, Ariz. | U | U | U | U | U | U | U |
| Detroit, Mich. | 178 | 106 | 40 | 18 | 7 | 7 | 11 | Pueblo, Colo. | 26 | 16 | 9 | 1 | - | - | 5 |
| Evansville, Ind. | 45 | 36 | 3 | 4 | 1 | 1 | 3 | Salt Lake City, Utah | 107 | 68 | 19 | 10 | 5 | 5 | 12 |
| Fort Wayne, Ind. | 53 | 37 | 10 | 4 | 2 | - | 4 | Tucson, Ariz. | 123 | 93 | 22 | 5 | 2 | 1 | 9 |
| Gary, Ind. | 15 | 6 | 6 | 1 | 2 | - | - | PACIFIC | 1,034 | 726 | 209 | 63 | 16 | 20 | 88 |
| Grand Rapids, Mich. | 79 | 59 | 10 | 2 | 5 | 3 | 5 | Berkeley, Calif. | 13 | 6 | 5 | - | - | 2 | - |
| Indianapolis, Ind. | 206 | 144 | 36 | 13 | 7 | 6 | 12 | Fresno, Calif. | 76 | 45 | 20 | 8 | 2 | 1 | 6 |
| Lansing, Mich. | 46 | 36 | 6 | 2 | 1 | 1 | 4 | Glendale, Calif. | U | U | U | U | U | U | U |
| Milwaukee, Wis. | 148 | 102 | 34 | 7 | 1 | 4 | 11 | Honolulu, Hawaii | 84 | 59 | 16 | 5 | - | 4 | 2 |
| Peoria, Ill. | 73 | 54 | 14 | 3 | 2 | - | 3 | Long Beach, Calif. | 67 | 56 | 7 | 3 | 1 | - | 9 |
| Rockford, Ill. | 67 | 55 | 7 | 1 | 1 | 3 | 4 | Los Angeles, Calif. | U | U | U | U | U | U | U |
| South Bend, Ind. | 45 | 34 | 7 | 2 | 2 | - | 1 | Pasadena, Calif. | 27 | 21 | 3 | 3 | - | - | 5 |
| Toledo, Ohio | 108 | 81 | 21 | 3 | 1 | 2 | 6 | Portland, Oreg. | 108 | 76 | 23 | 4 | 4 | 1 | 2 |
| Youngstown, Ohio | 59 | 49 | 6 | 3 | - | 1 | 2 | Sacramento, Calif. | 158 | 108 | 41 | 6 | - | 3 | 14 |
| W.N. CENTRAL | 541 | 393 | 97 | 33 | 10 | 8 | 49 | San Diego, Calif. | 149 | 106 | 25 | 10 | 4 | 4 | 15 |
| Des Moines, Iowa | 77 | 62 | 12 | 2 | 1 | - | 11 | San Francisco, Calif. | U | U | U | U | U | U | U |
| Duluth, Minn. | 38 | 26 | 9 | 2 | 1 | - | 1 | San Jose, Calif. | 161 | 109 | 39 | 9 | 1 | 3 | 18 |
| Kansas City, Kans. | 49 | 28 | 12 | 7 | 2 | - | 3 | Santa Cruz, Calif. | 30 | 23 | 4 | 2 | 1 | - | 5 |
| Kansas City, Mo. | 74 | 57 | 12 | 5 | - | - | 4 | Seattle, Wash. | 113 | 78 | 21 | 11 | 2 | 1 | 7 |
| Lincoln, Nebr. | 40 | 31 | 4 | 3 | 1 | 1 | 5 | Spokane, Wash. | 48 | 39 | 5 | 2 | 1 | 1 | 5 |
| Minneapolis, Minn. | 51 | 35 | 12 | 2 | - | 2 | 8 | Tacoma, Wash. | U | U | U | U | U | U | U |
| Omaha, Nebr. | 78 | 56 | 15 | 4 | - | 3 | 11 | | | | | | | | |
| St. Louis, Mo. | U | U | U | U | U | U | U | | | | | | | | |
| St. Paul, Minn. | 49 | 39 | 8 | 1 | - | 1 | 4 | | | | | | | | |
| Wichita, Kans. | 85 | 59 | 13 | 7 | 5 | 1 | 2 | | | | | | | | |

U: Unavailable. - : No reported cases.

* Mortality data in this table are voluntarily reported from 122 cities in the United States, most of which have populations of $\geq 100,000$. A death is reported by the place of its occurrence and by the week that the death certificate was filed. Fetal deaths are not included.

† Pneumonia and influenza.

‡ Because of changes in reporting methods in this Pennsylvania city, these numbers are partial counts for the current week. Complete counts will be available in 4 to 6 weeks.

§ Total includes unknown ages.

All *MMWR* references are available on the Internet at <http://www.cdc.gov/mmwr>. Use the search function to find specific articles.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.


References to non-CDC sites on the Internet are provided as a service to *MMWR* readers and do not constitute or imply endorsement of these organizations or their programs by CDC or the U.S. Department of Health and Human Services. CDC is not responsible for the content of these sites. URL addresses listed in *MMWR* were current as of the date of publication.

The *Morbidity and Mortality Weekly Report (MMWR)* Series is prepared by the Centers for Disease Control and Prevention (CDC) and is available free of charge in electronic format and on a paid subscription basis for paper copy. To receive an electronic copy each week, send an e-mail message to listserv@listserv.cdc.gov. The body content should read *SUBscribe mmwr-toc*. Electronic copy also is available from CDC's World-Wide Web server at <http://www.cdc.gov/mmwr> or from CDC's file transfer protocol server at <ftp://ftp.cdc.gov/pub/publications/mmwr>. To subscribe for paper copy, contact Superintendent of Documents, U.S. Government Printing Office, Washington, DC 20402; telephone 202-512-1800.

Data in the weekly *MMWR* are provisional, based on weekly reports to CDC by state health departments. The reporting week concludes at close of business on Friday; compiled data on a national basis are officially released to the public on the following Friday. Address inquiries about the *MMWR* Series, including material to be considered for publication, to Editor, *MMWR* Series, Mailstop C-08, CDC, 1600 Clifton Rd., N.E., Atlanta, GA 30333; telephone 888-232-3228.

All material in the *MMWR* Series is in the public domain and may be used and reprinted without permission; citation as to source, however, is appreciated.

☆U.S. Government Printing Office: 2003-533-155/69077 Region IV



DEPARTMENT OF HEALTH AND HUMAN SERVICES
CENTERS FOR DISEASE CONTROL AND PREVENTION (CDC)
ATLANTA, GA 30333

OFFICIAL BUSINESS
PENALTY FOR PRIVATE USE \$300
RETURN SERVICE REQUESTED

0206 93036 N1633DS 0001
PROQUEST INFORMATION & LEARNING
PERIODICALS ACQUISITION
PO BOX 1346
ANN ARBOR MI 48106-1346

FIRST-CLASS MAIL
POSTAGE & FEES PAID
PHS/CDC
Permit No. G-284

